PHENYLEPHRINE HCl
CHLORPHENAMINE MALEATE

DISUDRIN®

5 mg / 1 mg per 5 mL Syrup
2.5 mg / 0.5 mg per mL Syrup (Oral Drops)

Nasal Decongestant / Antihistamine

FORMULATIONS
Each mL syrup (oral drops) contains:
Phenylephrine HCl ..... 2.5 mg
Chlorphenamine maleate ..... 0.5 mg

Each 5 mL (1 teaspoonful) syrup contains:
Phenylephrine HCl ..... 5 mg
Chlorphenamine maleate ..... 1 mg

PRODUCT DESCRIPTIONS
Phenylephrine HCl + Chlorphenamine maleate (Disudrin®) Syrup (Oral Drops) is an orange, clear syrup with orange taste.
Phenylephrine HCl + Chlorphenamine maleate (Disudrin®) Syrup is an orange, clear syrup with fruity taste.

WHAT IS IN THE MEDICINE?
These medicines contain Phenylephrine HCl and Chlorphenamine maleate.
Phenylephrine HCl, a nasal decongestant, clears obstructed air passages and nasal sinuses due to congestion making breathing easier. It also reduces postnasal drip.
Chlorphenamine maleate, an anti-allergy, relieves symptoms of allergy such as runny nose, sneezing, and itchy, watery eyes.

STRENGTH OF THE MEDICINE
See Formulations

WHAT IS THIS MEDICINE USED FOR?
These medicines are used for the relief of clogged nose, runny nose, itchy, watery eyes, and sneezing associated with the common cold, upper airway cough syndrome (postnasal drip syndrome), allergic rhinitis, and other minor respiratory tract infections. They also help decongest sinus openings and passages.

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?
These medicines should be taken orally (by mouth).

Age Group	Oral Dose of Phenylephrine HCl + Chlorphenamine maleate (Disudrin®) every 6 hours	
	2.5 mg / 0.5 mg per mL Syrup (Oral Drops)	**5 mg / 1 mg per 5 mL Syrup**
Children 1 to 3 months 4 to 6 months 7 to 12 months 1 to 2 years	 0.25 mL 0.5 mL 0.75 mL 1 mL	 - - - -
2 to 6 years 7 to 12 years Children 12 years and older and Adults	- - -	2.5 mL (1/2 teaspoonful) 5 mL (1 teaspoonful) 10 mL (2 teaspoonsful)
	Or, as recommended by a doctor.	

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?
- If you are allergic to any ingredient in the product
- If you have high blood pressure or severe heart disease unless recommended by a doctor
- If you are pregnant or breastfeeding

UNDESIRABLE EFFECTS
Phenylephrine HCl may cause tremors (muscle shaking), restlessness, anxiety (feeling of uneasiness), insomnia/sleeplessness, nervousness, dizziness, increased blood pressure, palpitation (rapid and irregular heart beat), arrhythmia (irregular heart beat), weakness, respiratory distress (noisy, congested breathing), and pallor (pale skin color).

Chlorphenamine maleate may cause sleepiness and drowsiness. Other undesirable effects include muscle weakness, gastrointestinal discomfort, nausea, vomiting, diarrhea or constipation, dryness of the mouth, nose, and throat, difficulty urinating, ringing in the ears, visual disturbance, blurred vision, insomnia/sleeplessness, tremors, nervousness, irritability, chest tightness, increased or decreased blood pressure, and headache.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Do not use this product together with sympathomimetic agents (e.g., Epinephrine, etc.) and general anesthetics (e.g., Halothane, etc.) because of the possibility for increased toxicity.
Concurrent administration with medicines for depression (e.g., Amitriptyline, Imipramine, Sertraline, Moclobemide, etc.) may result in hypertensive crisis (sudden, severe increase in blood pressure that can lead to stroke).
Taking an adrenergic-blocking agent (e.g., Phenothiazine drugs, Phentolamine, Propranolol, etc.) before the administration of this product may decrease the effect of phenylephrine.
Tell your doctor about other medicines the child is taking, especially other medicines for cough, cold or allergy.

WHAT SHOULD YOU DO IF YOU MISS A DOSE?

If you miss a dose, just take the next dose if still needed for the condition being treated, and the subsequent doses at the recommended time or schedule (i.e., every 6 hours).

Do not double the dose.

HOW SHOULD YOU KEEP THIS MEDICINE?

Keep the product out of reach and sight of children
Store at temperatures not exceeding 30°C

SIGNS AND SYMPTOMS OF OVERDOSAGE

Phenylephrine HCl

- Increased blood pressure
- Palpitation
- Vomiting
- Headache
- Paresthesia (sensation of tingling, pricking, or numbness of the skin)
- Cerebral hemorrhage (bleeding from a ruptured blood vessel in the brain)
- Seizures

Chlorphenamine maleate

- Extreme sleepiness or prolonged drowsiness
- Convulsion
- Tachycardia (fast heart rate)
- Weakness
- Agitation
- Coma
- Hallucination
- Irritability

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

If you have taken more than the recommended dosage, consult a doctor.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

- May cause drowsiness; patients should observe caution while driving or performing other tasks requiring alertness.
- Do not use after the expiry date on the label.

WHEN SHOULD YOU CONSULT YOUR DOCTOR?

Before taking this medication, tell your doctor if you have:

- High blood pressure or any type of heart problems
- Thyroid problems
- An enlarged prostate, bladder problems or difficulty urinating
- Glaucoma
- Diabetes

If symptoms do not improve or new symptoms appear

AVAILABILITY

Phenylephrine HCl + Chlorphenamine maleate (Disudrin®) 2.5 mg / 0.5 mg per mL Syrup (Oral Drops)
– in amber bottle of 10 mL
Phenylephrine HCl + Chlorphenamine maleate (Disudrin®) 5 mg / 1 mg per 5 mL Syrup
– in amber bottle of 60 mL & 120 mL

DATE OF FIRST AUTHORIZATION (DR-XY29955): March 2011
DATE OF FIRST AUTHORIZATION (DR-XY33744): March 2011

DATE OF REVISION OF PATIENT INFORMATION LEAFLET (PIL): April 2011

ADVERSE DRUG REACTION REPORTING STATEMENT

For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph **AND** Unilab at (+632) 858-1000 or productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

Manufactured by AMHERST LABORATORIES, INC.
UNILAB Pharma Campus, Barangay Mamplasan
Biñan, Laguna, Philippines
for **UNILAB, Inc.**
No. 66 United Street, Mandaluyong City, Metro Manila, Philippines

Trusted Quality Healthcare

REG. IPOPHIL