Ketorolac Trometamol Kortezor®

30 mg/mL solution for intramuscular (IM)/intravenous (IV) injection Non-steroidal anti-inflammatory drug / Analgesic / Antipyretic

FORMULATION
Each mL solution for injection contains
Ketorolac trometamol ..... 30 mg
Water for injection ..... q.s.

PRODUCT DESCRIPTION
Kortezor® (ketorolac trometamol)® is a clear, slightly yellow, sterile, aqueous solution, free from extraneous matter and filled as a 2 mL ampoule with blue line.

PHARMACODYNAMICS
Ketorolac trometamol exhibits anti-inflammatory, analgesic and antipyretic activities by inhibiting cyclooxygenase (COX) 1 and COX 2A. Ketorolac trometamol is a potent inhibitor of prostaglandin synthesis. Ketorolac trometamol has no opioid agonist or antagonist properties. It does not suppress the respiratory system and is not addictive.

The greatest difference between large and small doses of ketorolac trometamol is in the duration of analgesia.

PHARMACOKINETICS
After IM administration of 10 or 30 mg doses of ketorolac trometamol injection in healthy adults, time to peak plasma concentrations is achieved at approximately half-life of about 45 to 50 minutes. In single dose studies, there is a reduction in clearance and an increase in terminal half-life (see table below).

The drug undergoes biotransformation in the liver to form p-hydroxyketorolac. The hydroxy metabolite exhibits limited pharmacologic activity. Having less than 20% or 1% of the anti-inflammatory or analgesic potency, respectively, of the parent drug. Ketorolac also undergoes conjugation with glucuronic acid. The drug and its metabolites are excreted almost completely by the kidneys, about 91.4% of the dose is recovered in the urine, and the remainder in the feces.

The primary mode of excretion of ketorolac trometamol and its metabolites (conjugates and the p-hydroxy metabolite) is in the urine (92% within 2 days of dosing). Fecal excretion accounts for less than 6% within 3 days. Renal clearance of ketorolac is rapid (in the order of 0.12 to 0.24 mL/min/kg), ketorolac glucuronide (25%-20%), p-hydroxyketorolac (17 % -12%), and unidentified polar metabolites (5-7%).

The Influence of Age, Liver, and Kidney Function on the Clearance and Terminal Half-Life of Ketorolac in Adult Populations[1]

PATIENT GROUP	TOTAL CLEARANCE (L/hr/kg) Mean (Range)	TERMINAL HALF- LIFE (hours) Mean (Range)
Normal Subjects	0.029 (0.017 to 0.050)	5.3 (3.5 to 9.2)
Healthy elderly subjects	0.021 (0.013 to 0.030)	7.3 (5.6 to 9.1)
Patients with hepatic dysfunction	0.030 (0.015 to 0.060)	5.4 (2.2 to 8.9)
Patients with renal impairment (creatinine= 1.9 to 9 mg/dL)	0.0017 (0.0006 to 0.0030)	10.3 (5.8 to 19.2)
Renal dialysis patients	0.016 (0.003 to 0.036)	13.8 (8 to 36)

[1]Estimated from 30 mg single IM doses of ketorolac trometamol

INDICATION
Short term (5 days) management of moderate to severe acute postoperative pain.

DOSAGE AND ADMINISTRATION
Ketorolac trometamol should be given the lowest effective dose of ketorolac should be used for the shortest possible time. Kortezor® (ketorolac trometamol)® is for intramuscular (IM) or intravenous (IV) use and the frequency should be adjusted to each individual patient needs.

General Dosing Recommendation
- Administered only IV or IM ketorolac should not exceed 2 days (see **Bowel Warnings**).
- Ketorolac is not for topical use.
- Ketorolac ampules should not be used for epidural or spinal administration.
- Ketorolac should not be mixed in the same syringe with morphine.
- Analgesic effects begin to appear about 30 minutes and peak in 1 to 2 hours after IV or IM administration.
- The analgesic effect may be maintained up to 6 to 8 hours.
- The total dose of ketorolac trometamol injection administered by any route or the injection rate for 15 to 30 seconds for intravenous administration.
- Strict aseptic technique should be observed when drawing up the contents of the ampule. As with all parenteral drug products, ketorolac trometamol injection should be inspected visually for particulate matter and discoloration prior to administration.
- Each ampule is for single use only. The solution should be used immediately after opening. Any unused contents should be discarded.
- For patients converted from parenteral to oral ketorolac, the total dose should not exceed 90 mg a 30 mg for
- Patients should be instructed that oral ketorolac is not to be used while receiving IM or IV ketorolac.
- Patients should be counseled that oral ketorolac is not to be used while receiving IM or IV ketorolac.
- Opioid analgesics (e.g., morphine, pethidine) may be used concomitantly, and may be required for optimal
- Concomitant use of ketorolac trometamol and probenecid is not recommended. Probenecid significantly
- Ketorolac trometamol should be used with caution if another nonsteroidal anti-inflammatory drug is
- Concomitant use of ketorolac trometamol with nephrotoxic drugs should be avoided unless the potential adverse effects should still be considered.

Recommended Ketorolac Adult IV/IM Dose:
- Initially, 10 mg followed by 10 to 30 mg every 4 to 6 hours, as needed
- For the initial postoperative period, ketorolac may be given as a flat dose on a every two hours if needed
- **Maximum Dose:** 90 mg/day

Special Populations

Geriatric (>65 years old):	A single dose with the lower end of the dosage range is recommended, the initial
Maximum Dose:	60 mg/day
Maximum Dose:	90 mg/day

Hepatic Impairment: Ketorolac is contraindicated in moderate to severe acute postoperative pain.

Renal Impairment: Ketorolac is contraindicated in moderate to severe renal impairment, reduce dosage in lower renal impairment (not to exceed 60 mg/day for IV IM).

If you are prescribed by a physician:

CONTRAINDICATIONS
- Known hypersensitivity to ketorolac trometamol or other ingredients in the product
- Patients with a history of asthma, urticaria, or allergic-type reactions after taking (prostaglandin) or perforation, or a history of documented ulceration or a documented active disease of the GI system.
- Patients with complete or partial syndrome of nasal polyps, or in whom asthma, anaphylactic, urticaria,
- Patients with suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis,
- Patients who have been recently (within 48 hours) been treated with aspirin or other NSAIDs. Severe, sometimes fatal, anaphylactoid reactions to NSAIDs have been reported in such patients.
- Asa prophylactic analgesics before surgery and in the systemically
- For nonsteroidal anti-inflammatory drugs and administration due to its alcohol content
- Significant hepatic impairment, active liver disease or severe liver failure.
- Patients with renal impairment (creatinine clearance less than 30 mL/min/1.73 m2), patients at risk of
- renal failure because of volume depletion or other factors.
- Patients with suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis, or patients at high risk of bleeding.
- Patients with severe heart failure.
- Patients in labor or delivery, as it may increase the risk of bleeding.
- Patients receiving concomitant aspirin or NSAIDs because of the increase of the cumulative risk of serious NSAID-related adverse effects.
- Patients who are breastfeeding or intended protocol, perioperative or lithium salts
- In patients not undergoing elective surgery with inadequate and low dose hemostasis (2.500 to 5.000 units every 12 hours).
- Pregnancy, labor, delivery, or breastfeeding
- Children below 16 years old
- **Contraindicated in the treatment of perioperative pain in the setting of coronary artery by-pass graft (CABG) surgery**

WARNINGS AND PRECAUTIONS

Treatment with Kortezor® should not exceed 2 days because adverse events may increase with prolonged use. There is limited experience with dosing for longer periods since often two days majority of patients no longer require parenteral administration of ketorolac trometamol. The total duration of combined IM and oral treatment should not exceed 5 days.

Cardiovascular Risk
- NSAIDs may cause an increased risk of serious cardiovascular thrombotic events, myocardial infarction and stroke which can be fatal. This risk may occur as early as the first use. The risk may increase with duration of use. Patients with cardiovascular disease or risk factors for cardiovascular disease may be at greater risk.

Gastrointestinal Risk
- NSAIDs, including ketorolac, cause an increased risk of serious GI adverse events including bleeding, ulceration and perforation of the stomach or intestine, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious GI events.

CARDIOVASCULAR EFFECTS (see Contraindications)

Cardiovascular Thrombotic Events
- Clinical trials of several COX-2 selective and NSAIDs of serious arteriothrombotic events (e.g. chest pain, shortness of breath, weakness, slurring of speech), which can be fatal. Patients risk factors for CV events may be estimated to see a physician immediately in case of such an event.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID does increase the risk of serious GI events.

Hypertension
- NSAIDs can lead to the onset of new hypertension or worsening of pre-existing hypertension, either of which may contribute to the increased incidence of cardiovascular events. Patients taking thiazides or loop diuretics may have impaired response to these therapies when taking NSAIDs. NSAIDs should be used with caution in treating elderly patients with hypertension. Blood pressure should be monitored closely during the initiation of NSAID treatment and throughout the course of therapy.

Congestive Heart Failure and Edema
- Fluid retention and edema have been observed in some patients taking NSAIDs. Ketorolac should be used with caution in patients with fluid retention or heart failure.

NSAIDs have a potential risk of hyperkalemia, particularly in patients with conditions such as diabetes mellitus or renal failure, elderly patients, or in patients receiving concomitant therapy with beta-adrenergic blockers, angiotensin converting enzyme inhibitors or some diuretics.

Renal Effects and Hyperkalemia
- NSAIDs, including ketorolac, can cause an increased risk of serious GI adverse events including bleeding, ulceration and perforation of the stomach or intestine, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients are at greater risk for serious GI events.

Anaphylactoid Reactions
- NSAIDs should be prescribed with extreme caution in those with a history of ulcer disease or GI bleeding since these patients have a significantly greater risk of developing a GI bleed or perforation during therapy with NSAIDs. Ketorolac should be a factor. Other factors that increase the risk for GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status.

Phenytoin and patients should remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. This should include discontinuation of the NSAID until a serious GI adverse event is ruled out. Patients should be informed of the risk of NSAIDs should be considered in patients taking aspirin for prophylaxis.

Close medical surveillance and caution should be exercised in patients with ulcerative colitis or with Crohn's disease as these conditions may be exacerbated.

GENITOURINARY EFFECTS

Some NSAIDs are associated with serious urinary symptoms (bladder pain, dysuria, urinary frequency), hematuria or renal papillary necrosis. Patients who occur at greater risk of developing these events with NSAIDs. Discontinuation of ketorolac trometamol therapy usually results in prompt recovery to the pre-treatment state. Use of ketorolac trometamol investigations or treatments are carried out.

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a NSAID may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and angiotensin-converting enzyme (ACE) inhibitors, and those with extracellular volume depletion. NSAID therapy is usually followed by recovery to the pre-treatment state.

As with other drugs that inhibit prostaglandin synthesis elevations of serum urea creatinine and potassium have been reported with ketorolac and may occur after one dose.

Patients with Impaired Renal Function
Ketorolac is CONTRAINDICATED in patients with severe renal disease (see **Contraindications**). Patients with known renal disease should receive no more than a single dose of ketorolac and be monitored closely for changes in renal function.

ANAPHYLACTOID REACTIONS (see Contraindications)
Anaphylactoid reactions may occur in patients without known prior exposure to NSAIDs. Ketorolac should not be given to patients with the aspirin triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after ingestion of or with NSAIDs. Ketorolac should be administered with caution in patients with pre-existing asthma and emergency help should be sought in cases where an anaphylactoid reaction occurs.

SKIN REACTIONS
NSAIDs, including ketorolac, can cause serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson syndrome (SJS), and toxic epidermal necrosis (TEN) which can be fatal. These serious events may occur without warning. Patients should be informed about the signs and symptoms of serious skin manifestations and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

HEPATIC EFFECTS
Borderline elevations of one or more liver tests may occur in up to 15% of patients taking NSAIDs including ketorolac. These laboratory abnormalities may progress, may remain unchanged, or may be transient with continuing therapy. In patients with normal liver function tests and with relatively healthy livers, fatal hepatitis associated with NSAIDs has occurred in about 1 in 2000 to 1 in 4000 patients. Ketorolac should be discontinued if abnormal liver tests persist or worsen or if clinical signs and symptoms consistent with liver disease develop or if systemic manifestations occur (e.g., eosinophilia, rash).

HEMATOLOGICAL EFFECTS
Anemia is sometimes seen in patients receiving NSAIDs including ketorolac. This may be due to fluid retention, occult or gross GI bleeding, or an incomplete picture of erythropoiesis. Patients on long-term treatment with NSAIDs, including ketorolac, should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia.

Ketorolac inhibits platelet aggregation and has been shown to prolong bleeding time in patients with normal bleeding function, bleeding times were varied, but not outside the normal range of two to eleven minutes. Ketorolac has prolonged effects from aspirin, platelet function returns to normal within 24 to 48 hours after discontinuation of ketorolac.

Postoperative wound hemorrhage has been reported in association with the immediate post-operative use of ketorolac. Therefore, ketorolac should not be used in patients who have operations with a high risk of hemostasis or incomplete hemostasis. Caution should be used when elective hemostasis is critical (e.g., in cosmetic or dry-case surgery).

Hematoma and other signs of wound hemorrhage and symptoms have been reported with the use of ketorolac. Physician should be alert for the possibility of hematoma formation in patients undergoing major surgery especially if they are receiving concomitant therapy with anticoagulants (see **Contraindications**).

ASEPTIC MENINGITIS
Aseptic meningitis with some NSAIDs, the symptoms of aseptic meningitis (stiff neck, severe headaches, nausea and vomiting, fever or clouding of consciousness) have been observed. Patients with autoimmune disorders (systemic lupus erythematosus, mixed connective tissue disease) seem to be pre-disposed. Therefore, in such patients, the physician must be vigilant to the development of this complication.

INTERACTION
As with other NSAIDs, ketorolac trometamol injection may mask the usual signs of infection.

OPHTHALMOLOGIC EFFECTS
Blurred vision and other eye disorders have been reported with the use of ketorolac trometamol injection and other NSAIDs. If such disturbances occur patients should undergo an ophthalmologic examination (see **Contraindications**).

CONCOMITANT DISEASES (see Contraindications)
Ketorolac should be used very cautiously and with close medical supervision in patients who may be adversely affected by changes in renal function. Such patients include those with renal disease, congestive heart failure, liver dysfunction, those taking diuretics and those who are undergoing major surgery requiring renal support. Patients who are dehydrated should be rehydrated prior to the patient, since the drug may inhibit platelet function and hemorrhage is possible.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE
Dizziness, drowsiness, visual disturbance, headache, vertigo, insomnia and depression have been reported with ketorolac use. If patients experience these or other similar adverse effects, they should not drive or operate machinery.

INTERACTIONS WITH OTHER MEDICAMENTS

CONCOMITANT DRUGS	EFFECTS
ACE Inhibitors, Angiotensin Receptor Antagonists	Decreased hypotensive effect and increased risk of renal impairment, particularly in volume-depleted patients
Anticoagulants, Thrombolytics (e.g., warfarin, heparin)	CONTRAINDICATED: increased risk of GI adverse events such as ulceration and bleeding and may enhance anticoagulant effects
Antiinvestinals (e.g., phenytoin, carbamazepine)	Sporadic cases of seizure have been reported with concomitant use
Aspirin, NSAIDs	CONTRAINDICATED: increased side effects and hemorrhage
Cholestyramine, Tamsulosin	Increased risk of nephrotoxicity
Corticosteroids	Increased risk of gastrointestinal bleeding and ulceration
Digoxin (e.g., furosemide, thiazides)	NSAIDs may exacerbate cardiac failure, reduce glomerular filtration rate (GFR) and increase plasma cardiac glycoside levels. Reduced diuretic effects and increased nephrotoxicity.
Lithium	CONTRAINDICATED: ketorolac reduces excretion of lithium (increased risk of toxicity)
Methotrexate	NSAIDs may reduce the excretion of methotrexate (increased risk of toxicity). Caution should be exercised when ketorolac and methotrexate is given concomitantly.
Nondepolarizing Muscle Relaxants	May potentiate effects of relaxants, resulting in apnea
Pentoxifylline (oxpentifylline)	CONTRAINDICATED: increased risk of bleeding when ketorolac is given with pentoxifylline
Probenecid	CONTRAINDICATED: decreased clearance and increased plasma concentration, total AUC, and half-life of ketorolac
Psychoactive drugs (e.g., fluoxetine, thiothixene, alprazolam)	Concomitant administration produces hallucination.
Quinolones	Possible increased risk of convulsions
Selective Serotonin Reuptake Inhibitors	Increased risk of GI bleeding
Zidovudine	Increase the risk of hematologic toxicity

Tell your doctor if you are taking other medicines, including vitamins, supplements, and herbal medicines

TREATMENT ON USAGE FOR HIGH RISK GROUPS

Pregnancy
Pregnancy Category C: Ketorolac trometamol is CONTRAINDICATED during pregnancy, labor or delivery. Avoid ketorolac use during pregnancy (particularly late pregnancy) because of the known effects of NSAIDs on the fetal cardiovascular system (closure of ductus arteriosus). Ketorolac trometamol injection should be considered.

Labor and Delivery
Ketorolac should not be used during labor and delivery because it may adversely affect fetal circulation and inhibit uterine contractions thereby its effect on prostaglandin synthesis. This may increase the risk of uterine hemorrhage. Uterine contractions may be inhibited by ketorolac, consequently the use or administration of prostaglandin synthesis. NSAIDs may increase the onset of normal delivery.

There may be increased bleeding tendency in both mother and child.

Lactation
Ketorolac has been detected in human milk at low concentrations and is CONTRAINDICATED in breastfeeding women (see **Contraindications**).

Fertility
The use of ketorolac trometamol injection, as with any drug known to inhibit COX1/prostaglandin synthesis, may impair fertility and is not recommended in women who have difficulty conceiving or are undergoing investigations of infertility. Ketorolac trometamol should be discontinued.

Children (< 16 years old)
The safety and effectiveness of ketorolac in children have not been established (see **Contraindications**).

Elderly (> 65 years old)
Elderly patients are more susceptible to a variety of adverse reactions from NSAIDs, the incidence of these adverse reactions increases with age. Elderly patients are at greater risk for serious GI events. Ketorolac should be administered with caution to the elderly as NSAIDs' clearance and volume of distribution decrease (see **Dosage and Administration**).

Renal Impairment
Ketorolac and its metabolites are eliminated by the kidneys, which in patients with reduced creatinine clearance will result in diminished plasma clearance of the drug (see **Contraindications and Warnings and Precautions**).

UNDESIRABLE EFFECTS
Most frequently reported adverse effects with NSAIDs are GI, of which peptic ulcer, with or without bleeding, in the most severe. If patients have recurrent symptoms.

The most frequently reported adverse effects (≥ 1) 72%) with ketorolac trometamol are somnolence, nausea, headache, vomiting, injection site pain, sweating, dizziness, and asthenia.

Less common adverse effects (< 1% of patients) include:

Blood and Lymphatic System: Anemia, agranulocytosis, aplastic anemia, bleeding at the operative site (rarely requiring blood transfusion), coagulation disorders, hemolytic anemia, leukopenia, purpura, increased bleeding time, leukocytosis, neutropenia, pancytosis, thrombocytopenia

Immune System: Anaphylactoid reactions, anaphylaxis, angioedema, asthma, hypotension, bronchospasm,flushing laryngeal/laryngeal edema, rash

Metabolism and Nutrition: Anorexia, edema, excessive thirst, hyperkalemia, hyponatremia, hyponatremia, hyponatremia, increased appetite, weight gain

Psychiatric: Abnormal dreams, abnormal thinking, anxiety, depression, difficulty in concentration, euphoria, hallucinations, insomnia, nervousness, paresthesia, psychosis, somnolence, tremor, vertigo

Nervous System: Aseptic meningitis, asthma, convulsions, extrapyramidal manifestations, functional disorders, hyperkinesia, hypertonia, hypesthesia, hypokinesia, migraine, paresthesia, vertigo

Special senses: Abnormal taste, blurred vision, eye irritation, hearing loss, optic neuritis, retinal hemorrhage, tinnitus, visual disturbance (e.g., blurred vision)

Cardiovascular: Bradycardia, cardiac failure, chest pain, flushing, hematoma, hemorrhage, hypertension, hypotension, pallor, palpitations, peripheral edema, postural hypotension, tachycardia, thrombosis, vasculitis, ventricular tachycardia

Respiratory, Thoracic and Mediastinal: Asthma, dyspnoea, hiccup, increased cough, pulmonary edema, rhinitis

Gastrointestinal: Abdominal pain/discomfort, anorexia, constipation, diarrhea, dry mouth, dyspepsia, eructation, esophagitis, excessive thirst, flatulence, gastritis, gastrointestinal bleeding, gastrointestinal ulcer, gastrointestinal perforation, hematemesis, hemorrhoids, increased appetite, indigestion, melena, pancreatitis, peptic ulcer, stomatitis, thirst, vomiting, ulcerative stomatitis, abdominal distention

Hepatic: Cholestatic jaundice, hepatitis, liver failure, liver function abnormalities

Renal and Genitourinary: Acute renal failure, dysuria, elevated BUN, elevated creatinine (e.g., 1.6 and 10.7% of 1,at% syndrome), escherichia, erythema multiforme, exfoliative dermatitis, pruritus, rash (may be maculopapular), skin disorder, skin photosensitivity, subcutaneous hematoma, urticaria

Musculoskeletal: Back pain, myalgia, neck pain, tendinitis

General Disorders and Administration Site Conditions: Chills, fatigue, fever, infections, injection site reaction or pain, malaise, pain, tingling at the injection site

Urogenital/Reproductive: Acute renal failure, dysuria, flank pain (with or without hematuria and/or azotemia), glomerular nephritis, hematuria, hyperkalemia, hyperuricemia, hypocalcemia, hypokalemia, hyponatremia, increased BUN, increased creatinine, interstitial nephritis, oliguria, polyuria, proteinuria, renal papillary necrosis, urinary retention, vaginitis

Investigations: Abnormal liver function tests, increased blood urea nitrogen (BUN) and creatinine, prolonged bleeding time

OVERDOSE AND TREATMENT
There is no specific antidote. 360 mg given 90% over an eight hour interval for five consecutive days caused abdominal pain and psychic ulcer which healed after discontinuation of dosing. There have been reports of metabolic acidosis following intentional overdosage.

Symptomatic and supportive care should be given to patients following an NONID overdose. There are no specific antidotes

Forced diuresis appears to be ineffective in removing the drug from circulation.

STORAGE CONDITION
Store out of sight and reach of children.
Store at temperatures not exceeding 30°C. Protect from light

AVAILABILITY
Ketorolac Trometamol (KORTEZOR®), 30mg/mL ampule Solution for Injection (box of 10 ampules)

CAUTION: Fraudu, Drugs, Overuse, and Counterfeits Act prohibits dispensing without prescription.

Manufactured by MARCELINO PHARMACEUTICALS
Sta. Rosa Tagaytay Road, Dasmariñas
Sta. Rosa City, Laguna, Philippines
Tel. 046 419 0888
No. 888 Andres Street, Mandaluyong City
Metro Manila, Philippines

Trusted Quality Health Care

Date of First Authorization : DR-XY22028-30 December 2003

P30000002160
Date of Revision : June 2018

Reg. IPOPHIL