Sodium chloride

NasoClear®

0.65% Saline Solution
Nasal Spray

FORMULATION

Each mL contains:
Sodium chloride ..... 6.5 mg
Benzalkonium chloride (as preservative) 0.02%

PRODUCT DESCRIPTION

Sodium chloride (NasoClear®) 0.65% Saline Solution Nasal Spray is a clear, colorless solution, free from foreign matter, supplied in a white plastic bottle with plastic cap (nasal atomizer).

WHAT IS IN THE MEDICINE?

This medicine contains sodium chloride solution, a salt solution that treats dry or irritated nasal passages. It works by rinsing and moisturizing the nostril.

Sodium chloride nasal spray is used to treat dryness inside the nose (nasal passages). It helps add moisture inside the nose to dissolve and soften thick crusty mucus and provide relief for dry, crusted, irritated, and inflamed nasal membranes due to colds, allergies, low humidity, heat, environmental factors, air travel, use of steroid nasal spray, overuse of topical nasal decongestants and other nasal solutions, chronic sinusitis, or rebound sinus reacting from smog. It helps relieve stuffiness and makes breathing easier.

For infants and young children with stuffy nose who cannot blow their noses, use of the product makes the mucus easier to remove with a nasal bulb syringe.

STRENGTH OF THE MEDICINE

See Formulation

WHAT IS THIS MEDICINE USED FOR?

This medicine is used to help provide relief from:

- Nasal congestion
- Cold or flu symptoms
- Nasal irritation due to pollen, dust mites, mold, and pollution
- Nasal allergies
- Upper airway cough syndrome (postnasal drip)

HOW MUCH AND HOW OFTEN SHOULD YOU USE THIS MEDICINE?

Children and Adults: Apply 2-4 sprays in each nostril 3 times a day, or, as directed by a doctor. May use a cotton bud to remove excess mucus. Use clean cotton to wipe the tip of the dropper after each use.

WHEN SHOULD YOU NOT TAKE THIS MEDICINE?

- If you are allergic to any ingredient in the product.

UNDESIRABLE EFFECTS

This product does not contain any medication and undesirable effects usually do not occur with use of this product. However, if the inside of your nose is very dry and irritated, stinging may occur. This product may also induce transient sneezing.

WHAT OTHER MEDICINE OR FOOD SHOULD BE AVOIDED WHILE TAKING THIS MEDICINE?

Tell your doctor about other medicines you are taking, especially other medicines for cough, cold or allergy.

WHAT SHOULD YOU DO WHEN YOU MISS A DOSE?

Just apply the missed dose if still needed for the particular condition being treated and then continue with your regular dosing schedule.

HOW SHOULD YOU KEEP THIS MEDICINE?

Keep the product out of reach and sight of children.
Store at temperatures not exceeding 30°C.

SIGNS AND SYMPTOMS OF OVERDOSAGE

There have been no reports of overdose with this product.

WHAT TO DO WHEN YOU HAVE TAKEN MORE THAN THE RECOMMENDED DOSAGE?

If you have used more than the recommended dosage, consult a doctor.

CARE THAT SHOULD BE TAKEN WHEN TAKING THIS MEDICINE

Do not share this product with others because of the possible increased risk of infection.
Do not use after the expiry date on the label.

WHEN SHOULD YOU CONSULT A DOCTOR?

If your condition persists or worsens.

AVAILABILITY

White Plastic bottle x 30 mL with plastic cap (Box of 1's)
DATE OF REVISION OF PATIENT INFORMATION LEAFLET (PIL) 06/2016

Manufactured by Hizon Laboratories, Inc.
Assumption Road, Sumulong Highway
Antipolo City, Philippines
Distributed by United Laboratories, Inc.
No. 66 United Street, Mandaluyong City, Metro Manila, Philippines
Under Authority by
UNILAB, Inc.
No. 66 United Street, Mandaluyong City, Metro Manila, Philippines

UNILAB

HLIPIN0021040402

Reg. IPOPHIL