Zoledronic Acid Osteomet

4 mg/5 mL Concentrate Solution for Intravenous Infusion
Bisphosphonate

FORMULATION
Each 5 mL vial contains:
Zoledronic acid (as monohydrate) ..... 4 mg

PHARMACOLOGIC CATEGORY

Bisphosphonate

PRODUCT DESCRIPTION

Zoledronic acid is a clear and colorless concentrated solution intended to be diluted prior to intravenous infusion.

CLINICAL PHARMACOLOGY

Pharmacodynamics
Zoledronic acid is an aminobisphosphonate whose main action is to inhibit bone resorption. The drug has a high affinity for mineralized bone but its precise mode of action in the inhibition of bone resorption is related to osteoclast activity. In animal studies, zoledronic acid clearly inhibits osteoclast activity by affecting a variety of intracellular processes necessary for osteoclast function. It induces apoptosis (programmed cell death) of osteoclasts, which are responsible for breakdown of bone tissue. It reduces the osteoclastic recruitment of mineralized bone and cartilage through its binding to bone. Zoledronic acid also inhibits the increased osteoclastic activity induced by tumor cells and prevents the bone destruction normally seen in tumor metastases.

Pharmacokinetics
Single and multiple 5- and 15-minute infusions of 2, 4, and 16 mg zoledronic acid in patients with bone metastases yielded the following pharmacokinetic data, which were found to be independent of dose. Following the start of the infusion, zoledronic acid plasma concentrations rapidly increased to a maximum (peak) value at the end of the infusion period. This was followed by a rapid decline to < 10% of peak after 4 hours and < 1% of peak after 24 hours with a subsequent prolonged period in which drug concentrations not exceeding 0.1% of peak prior to the second infusion of drug on day 28. There was no dose accumulation of zoledronic acid in plasma after multiple dose given every 28 days.

Zoledronic acid showed no affinity for the cellular components of the blood. Plasma protein binding in vitro was low (approximately 56%) and there was no evidence of saturation at these levels.

Zoledronic acid is not metabolized and is excreted unchanged via the kidney. Over the first 24 hours, 39 ± 16% of the administered dose is found in the urine, the rest being bound to bone or circulating bound to the plasma proteins. The terminal plasma elimination half-life is released very slowly, being more than 146 hours and determined by the kidney. The total body clearance is 5.04 ± 2.5 but and is independent of the dose. Renal clearance is 4.37 ± 2.5 but and accounts for 84% ± 16% of drug removal.

Increasing the infusion time from 5 to 15 minutes caused a 30% decrease in zoledronic acid concentration at the end of the infusion, but this had no significant effect on the plasma concentration-versus-time curve.

Renal Impairment
The renal clearance of zoledronic acid was correlated with creatinine clearance (Cl_{cr}), renal clearance representing 75 ± 33% of the Cl_{cr}, which showed a mean of 84 ± 29 mL/min/as (range 22 to 143 mL/min) in patients with cancer and bone metastases. Patients with mild renal impairment (Cl_{cr} 61-90 mL/min/as) cleared mean zoledronic acid 1.3 times more than patients with normal renal function (Cl_{cr} > 90 mL/min/as). The clearance of zoledronic acid in patients with severe renal insufficiency (Cl_{cr} < 30 mL/min/as). The use of zoledronic acid is not recommended in patients with severe renal impairment.

Hypercalcemia or Hepatic Insufficiency
No pharmacokinetic studies of zoledronic acid are available in patients with hypercalcemia or in patients with hepatic insufficiency.

Effects of Gender, Age and Race
Pharmacokinetic studies in patients with bone metastases showed that gender, race, age (between 38 and 84 years) and body weight had no significant clinical relevance on clearance.

INDICATIONS

- For the prevention of skeletal related events (pathological fractures, spinal compression, radiation or surgery to bone, or tumor-induced hypercalcemia) in adult patients with advanced malignancies involving bone
- For the treatment of adult patients with tumor-induced hypercalcemia (TIH).

DOSAGE AND ADMINISTRATION

General Dosing Considerations

Aseptic technique must be followed during the preparation of the infusion. Visually inspect the solution for particulate matter or discoloration prior to use, whenever the solution and container permit.

This product is for intravenous infusion only.

This product is intended for single use only. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Dosing Recommendations

Prevention of skeletal related events in patients with advanced malignancies involving bone:
The dose given as an intravenous infusion lasting for no less than 15 minutes every 3 to 4 weeks. The concentrate must be diluted with 100 mL 0.9% w/v sodium chloride or 5% w/v glucose solution prior to administration.

Hypercalcemia of Malignancy (HCM) (albumin-corrected serum calcium > 12 mg/dL (3 mmol/L)):
The dose as a single dose intravenous infusion over not less than 15 minutes following standard rehydration procedures.

Hydration: Patients must be rehydrated in a well hydrated state prior to and following administration of zoledronic acid. Promptly initiate vigorous saline hydration (an integral part of hypercalcemia therapy) and ensure adequate hydration throughout the study period. Only throughout treatment. Mild or asymptomatic hypercalcemia may be treated with conservative measures, including hydration with normal saline solution. Diuretics should be administered after adequate rehydration is established and should not be used to induce hypovolemia in patients who have cardiac failure. Do not empty diuretic therapy prior to correction of hypovolemia.

Renal Impairment: Do not administer zoledronic acid to patients with severe renal impairment at initial treatment, retreatment with zoledronic acid 4 mg may be considered. It is recommended that serum creatinine be measured before each zoledronic acid dose and treatment. It should be withheld if renal function has deteriorated. Recommended guidelines for re-starting zoledronic acid therapy provide deterioration in serum creatinine by 0.5 mg/dL (44 µmol/L) over baseline in patients with normal baseline.

In addition, retreatment should be given only to those patients who can tolerate the standard recommended dose of 4 mg. A 3.5 fold to 4 fold rise in Cl_{cr} (i.e. from more than 400-450 mL/min/as chloride per day). Serum BUN and creatinine must be evaluated and possible deterioration in renal function after each dose of zoledronic acid. Treatment should be withheld if renal deterioration is observed.

Dosage Adjustment in Patients with Renal Impairment
The risks and benefits of treatment with zoledronic acid should be evaluated in patients with HCM who have severe renal impairment. In clinical trials, patients with serum creatinine > 400 micromol/L or > 4.5 mg/dL were excluded. Dosage adjustments are not necessary in patients with mild to moderate renal impairment, i.e., serum creatinine < 400 micromol/L or < 4.5 mg/L. Zoledronic acid should not be used in patients with severe renal impairment.

Bone Metastases of Solid Tumors and Osteolytic Lesions of Multiple Myeloma (for patients with creatinine clearance of 60 mL/min or less):

4 mg given as a single dose IV infusion over not less than 15 minutes every 3 to 4 weeks

In patients requiring antineoplastic therapy, zoledronic acid should be administered either before or after the therapy. The concentrate must be diluted with 100 mL 0.9% w/v sodium chloride.

Patients will be required to take oral supplementation of calcium (500 mg) and vitamin D (400 IU) daily. Due to the risk of hypocalcemia, patients with impaired renal function should be carefully monitored for signs and symptoms related to altered calcium and Vitamin D.

Dosage Adjustment in Patients with Renal Impairment
In patients with mild to moderate renal impairment (defined as creatinine clearance 30-60 mL/min/as or 0.5-1.4 mg/dL), no dosage adjustment is necessary. Prior to each dose of zoledronic acid, renal function should be evaluated in order to achieve the same level of safety as that observed in patients with creatinine clearance of 75 mL/min. Creatinine clearance (Cl_{cr}) is calculated using the Cockcroft-Gault formula:

$$\text{Male: } Cl_{cr} = \frac{(140 - \text{age (years)}) \times (\text{Actual body weight (kg)})}{72 \times \text{Serum creatinine (micromol/L)}}$$

Female: 0.85 x value calculated using the above formula

Reduced Dose for Patients with Baseline Creatinine Clearance	Cl_{cr} < 60 mL/min Reduction in Zoledronic Acid Dose (given every 3-4 weeks)
>60	0 mg
50-60	3.5 mg
40-49	3.3 mg
30-39	3.0 mg

Doses calculated assuming target Cl_{cr} of 0.85(mg/mL) / (L/hr) = 75 mL/min)

Serum creatinine should be measured before each zoledronic acid dose and treatment. It should be withheld if renal function has deteriorated. Re-starting zoledronic acid therapy should be based on previous dose:

- For patients with normal baseline serum creatinine (< 4 mg/dL), an increase of 0.5 mg/dL.
- For patients with abnormal baseline serum creatinine (> 4 mg/dL), an increase of 0.4 mg/dL.

Direction for dilution
Using a sterile syringe, withdraw the contents of zoledronic acid concentrated solution (entire content of 5 mL vial), and dilute immediately in 100 mL of **sterile 0.9% sodium chloride or 5% dextrose injection prior to administration.**

Preparation of Reduced Doses of *Zoledronic Acid* for Patients with Baseline Cl_{cr} < 60 mL/min
Reduced doses of zoledronic acid concentrate needed as follows:
4.4 mL for 3.5 mg dose
4.1 mL for 3.3 mg dose
3.8 mL for 3.0 mg dose
The withdrawn amount of concentrate must be diluted immediately in 100 mL of **sterile 0.9% sodium chloride solution or 5% glucose solution.**

*Note: Smaller infusion volumes (e.g., 50 mL) and rapid (over 5 minutes) IV infusion rates have been associated with an increased risk of renal adverse reactions. Do not use these dose and rate. Strict adherence to administration recommendations for *Zoledronic Acid* concentrate is recommended.*

Incompatibilities
Due to lack of data on incompatibilities, zoledronic acid concentrate is to be diluted with 0.9% sodium chloride solution or 5% glucose solution.

Do not mix zoledronic acid with calcium or other divalent cation containing infusion solutions, such as Lactated Ringer's Solution, and should be administered as a single intravenous solution separate from all other drugs.

No incompatibilities with zoledronic acid have been detected in studies performed with several types of commonly used medical devices. Zoledronic acid is compatible with polyvinylchloride containers, Duotrol-PE tubing and Viaflex-polyolefin IV pouches as tested in beforehand and glass bottles.

Special Precautions for Storage:
Diluted zoledronic acid solution with 100 mL 0.9% sodium chloride or 5% glucose solution is stable at 2°C to 8°C for 24 hours.

Do not store unlabeled concentrate in syringe, to avoid inadvertent injection.

The diluted solution of zoledronic acid should be used immediately. If not used immediately, in use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2°C to 8°C. The refrigerated solution should then be equilibrated to room temperature prior to administration.

CONTRAINDICATIONS

- Hypersensitivity to zoledronic acid, other bisphosphonates or to any components of the product
- Pregnant or breastfeeding women

WARNINGS AND PRECAUTIONS

General
Patients receiving zoledronic acid should be under the supervision of a physician experienced in the administration of chemotherapeutic agents.

Standard hypercalcemia-related metabolic parameters, such as serum levels of calcium, phosphate and magnesium, should be closely monitored following the initiation of therapy. Short-term supplemental therapy may be necessary if hypocalcemia occurs.

Local reactions occurred following intravenous administration of each dose of zoledronic acid. Lower doses of zoledronic acid are recommended in patients with bone metastases with mild to moderate renal impairment. Zoledronic acid should not be administered with any other potentially nephrotoxic drug. Patients must be monitored at the same dose as that given prior to administration of zoledronic acid. **Dosing Recommendations**

Renal Impairment
Before administering zoledronic acid, have been associated with renal toxicity manifested as deterioration of renal function, progression to renal failure and renal dialysis. Evaluation of renal function should be assessed to ensure that they are adequately hydrated. The use of drug therapy is not recommended. Laboratory evaluation of renal function after each dose. **Dosing Recommendations**

Renal Impairment
Before administering zoledronic acid, have been associated with renal toxicity manifested as deterioration of renal function, progression to renal failure and renal dialysis. Evaluation of renal function should be assessed to ensure that they are adequately hydrated. The use of drug therapy is not recommended. Laboratory evaluation of renal function after each dose. **Dosing Recommendations**

Risk factors predisposing patients to renal deterioration include dehydration, pre-existing renal impairment, chemotherapy, concomitant therapy with other nephrotoxic drugs, dosage, infusion volume and time. Zoledronic acid dose.

Transient increases in serum creatinine concentration may be greater in patients with impaired renal function.

Musculoskeletal Pain
Severe and occasionally incapacitating bone, joint and/or muscle pain have been reported infrequently in patients taking bisphosphonates. Such reports have been received after initiation of treatment. Recurrence of symptoms has been observed in some patients when re-challenged with the same medicine or another bisphosphonate.

Osteonecrosis of the Jaw (ONJ)
ONJ has been associated with the use of bisphosphonates. ONJ has been reported mostly in cancer patients treated with intravenous bisphosphonates, including zoledronic acid.

Risk factors for ONJ include cancer diagnosis, concomitant therapy (chemotherapy, angiogenesis therapy) or ONJ include steroid (oral bisphosphonates or numerous), maxillofacial pain, dental surgery (e.g., tooth extraction, bone teeth, osteomyelitis, loss of teeth, soft tissue disease). Dental recurrent or persistent soft tissue infection in the oral cavity and marked oral ulcer.

Do not use factors for ONJ. Before administering zoledronic acid, a dental examination with preventive dentistry is recommended. Patients with known risk factors for ONJ should avoid invasive dental procedures if possible. During treatment and after treatment for at the same dose as that given prior to administration of zoledronic acid.

Hydration and Electrolyte Monitoring
Before administering zoledronic acid, hypocalcemic patients should be assessed to assure that they are adequately hydrated. The use of drug therapy is not recommended. Use zoledronic acid with caution in patients with other forms of fluid overload. **Dosing Recommendations**

Renal Impairment
Before administering zoledronic acid, have been associated with renal toxicity manifested as deterioration of renal function, progression to renal failure and renal dialysis. Evaluation of renal function should be assessed to ensure that they are adequately hydrated. The use of drug therapy is not recommended. Laboratory evaluation of renal function after each dose. **Dosing Recommendations**

Risk factors predisposing patients to renal deterioration include dehydration, pre-existing renal impairment, chemotherapy, concomitant therapy with other nephrotoxic drugs, dosage, infusion volume and time. Zoledronic acid dose.

Transient increases in serum creatinine concentration may be greater in patients with impaired renal function.

Musculoskeletal Pain
Severe and occasionally incapacitating bone, joint and/or muscle pain have been reported infrequently in patients taking bisphosphonates. Such reports have been received after initiation of treatment. Recurrence of symptoms has been observed in some patients when re-challenged with the same medicine or another bisphosphonate.

Osteonecrosis of the Jaw (ONJ)
ONJ has been associated with the use of bisphosphonates. ONJ has been reported mostly in cancer patients treated with intravenous bisphosphonates, including zoledronic acid.

Risk factors for ONJ include cancer diagnosis, concomitant therapy (chemotherapy, angiogenesis therapy) or ONJ include steroid (oral bisphosphonates or numerous), maxillofacial pain, dental surgery (e.g., tooth extraction, bone teeth, osteomyelitis, loss of teeth, soft tissue disease). Dental recurrent or persistent soft tissue infection in the oral cavity and marked oral ulcer.

Do not use factors for ONJ. Before administering zoledronic acid, a dental examination with preventive dentistry is recommended. Patients with known risk factors for ONJ should avoid invasive dental procedures if possible. During treatment and after treatment for at the same dose as that given prior to administration of zoledronic acid.

Physical Fractures of the Femur
In bisphosphonate-treated patients, atypical, low-energy, or low trauma fractures of the femoral shaft have been reported. These fractures can occur anywhere in the femoral shaft from just below the lesser trochanter to just above the supracondylar flare. These transverse or short oblique fractures occur following minimal trauma to the femur.

Atypical femoral fractures most commonly occur with minimal or no trauma to the affected area. They may be bilateral and many patients experience prodromal pain preceding as fail, aching pain in the thigh or groin, often associated with imaging features of stress fracture, weeks to months before a complete fracture of the femoral shaft occurs. The combination therapy with glucocorticoid in zoledronic acid treated patients who have sustained a femoral shaft fracture. Poor healing of these fractures has also been reported.

Evaluate any patient with a history of bisphosphonate exposure who presents with thigh or groin pain for an incomplete femoral fracture. Discontinuation of bisphosphonate therapy should be considered, pending a definitive assessment, on an individual basis.

Patients with Fractures
There have been reports of bronchoconstriction in aspirin-sensitive patients receiving bisphosphonates. Use zoledronic acid with caution in aspirin-sensitive patients.

Ophthalmia
Zoledronic acid may cause ocular disturbances (conjunctivitis, uveitis, episcleritis, scleritis, and orbital inflammation). Patients with ocular events other than uncomplicated conjunctivitis should be promptly referred to an ophthalmologist. Dosing Recommendations and Administration

Effects on Ability to Drive or Use Machines
No studies on the effects on the ability to drive and/or dizziness. Patients receiving zoledronic acid should be advised to take precautions when driving a vehicle or using machines until redness or physical combination. It should be carefully considered whether it is advisable to drive or operate machinery under these circumstances.

DRUG INTERACTIONS

There have been no specific drug-drug interaction studies conducted with zoledronic acid. Since zoledronic acid is not metabolized by the liver, it is unlikely to alter the in vivo liver and microsome CYP450 enzymes, interactions resulting from displacement of highly protein-bound drugs. Zoledronic acid does not inhibit the drug metabolizing enzyme cytochrome P450 in vitro.

Zoledronic acid has been administered concomitantly with the following drugs without clinically apparent interactions: Anticancer agents, diuretics, antibiotics and analgesics.

Aminoglycosides: Concomitant administration may result in possible additive effect in lowering serum calcium for prolonged period of time.

Nephrotoxic Drugs (e.g., aminoglycosides, other antineoplastic agents, ASA, NSAIDs) or drugs that significantly impact renal function (e.g., diuretics, ACE inhibitors): Zoledronic acid should not be given with such medication administered close in time because of the increased risk of renal impairment.

Thalidomide: The risk of renal insufficiency may be increased when zoledronic acid is used concomitantly with thalidomide especially in patients with multiple myeloma.

STATEMENT ON USAGE FOR HIGH RISK GROUPS

Pregnancy Category D

Zoledronic acid should not be used during pregnancy. Although no data exists on fetal risk in humans, animal reproduction studies have revealed that the bone mass of pups is reduced over a period of years. This may theoretically cause serious skeletal malformation when a fetus is exposed to zoledronic acid.

Lactation
It is not known whether zoledronic acid is excreted in human milk. Since many drugs are excreted in human milk and because zoledronic acid binds to long term, zoledronic acid should not be used by breastfeeding women.

Use in Infants and Children
The safety and efficacy of zoledronic acid in children 1 to 17 years old have not been established.

Use in the Elderly
There are no age-related differences in zoledronic acid's adverse events profile. However, since decreased renal function occurs more commonly in the elderly, special care should be taken to monitor renal function.

Hepatic Impairment
The clinical data available for the use of zoledronic acid in patients with hepatic impairment is limited; no dosage recommendations may be given in this group.

ADVERSE EFFECTS

Zoledronic acid is generally well tolerated and adverse effects are usually mild and do not require discontinuance of the drug.

Body as a Whole: Fever and flu-like symptoms (usually at the start of treatment), fatigue, chills, pain, malaise and burning, progression of cancer, weakness

Cardiovascular: Hypotension (associated with syncope or circulatory collapse primarily in patients with risk factors), arrhythmia, tachycardia, bradycardia, atrial fibrillation, edema, and hypertension (including worsening)

CNS: Insomnia, sleep disorder/disturbance, somnolence, anxiety, depression, confusion, agitation, headache, dizziness, vertigo, asthenia, paresthesia, taste disturbance, hypoesthesia, hyperesthesia, tremor, convulsion

Gastrointestinal Effects: Abdominal pain, nausea, anorexia, dyspepsia, constipation, diarrhea, vomiting, dry mouth, stomatitis, gastritis, dysphagia, weight loss, gastrointestinal disorder, abdominal distention, dry mouth, mild abdominal pain

Hemic and Lymphatic System: Anemia, neutropenia, granulocytopenia, thrombocytopenia, leukopenia, lymphopenia

Hypersensitivity Reactions: Rare reports of allergic reactions including angioedema, very rare cases of anaphylactic reactions/shock

Infection: Herpes simplex

Injection Site Reactions: Local reactions at the infusion site (including redness or swelling, pain, irritation, induration, and rash)

Laboratory Abnormalities: Hypophosphatemia, hypomagnesemia, hyperkalemia, hypokalemia, hypernatremia, increased blood creatinine and blood urea, hypocalcemia

Musculoskeletal: Osteonecrosis of the jaw (see Warnings), Musculoskeletal pain (which may be severe and incapacitating), bone pain, myalgia, arthralgia, back pain, pain in the limb, muscle cramp, musculoskeletal pain, joint stiffness, ONJ, atypical subtrochanteric and diaphyseal femoral fractures

Ocular: Uveitis, scleritis, episcleritis, conjunctivitis, blurred vision, chills, and orbital inflammation

Renal and Urinary Disorders: Renal impairment, acute renal failure, renal deterioration, hematuria, proteinuria, urinary tract infection

Respiratory: Dyspnea, cough, upper respiratory tract infection, bronchoconstriction

Skin and Subcutaneous Tissue Disorders: Alopecia, dermatitis, hyperhidrosis, pruritus, urticaria, rash (including erythematous and macular rash)

OVERDOSAGE AND MANAGEMENT

There is limited information on acute zoledronic acid overdosage. Overdosage would likely result in clinical abnormalities, hypocalcemia and hypophosphatemia which may be corrected by intravenous administration of calcium gluconate, potassium or sodium phosphate, and magnesium sulfate, respectively.

In two patients who received zoledronic acid 32 mg over 5 minutes in clinical trials, neither experienced any clinical or laboratory toxicity.

A female patient received a single 48 mg dose of zoledronic acid in error in an open label study of zoledronic acid and presented with decreased serum ionized magnesium (0.4°C), which resolved after treatment, two days after the overdose. All other evaluations were normal, and the patient was discharged seven days after the overdose.

Carefully monitor patients who have received doses higher than those recommended since renal function impairment including renal failure and serum electrolyte (including calcium, phosphorus and magnesium) abnormalities have been observed.

AVAILABILITY

Zoledronic Acid (Osteomet): 4 mg/5 mL vial - 5 mL clear PVC glass vial of 100 mg/5 mL cap (Box of 1's).

CAUTION

Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

For suspected adverse drug reaction, send medical attention immediately and report to the FDA at www.fda.gov.ph AND Unilab at +6222 809-1000 or productsafety@unilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

STORAGE

Store at temperature not exceeding 30°C.

Keep out of reach and sight of children.

Store in original packaging.

Manufactured by Pharmbiotest Sp.z.o.o.
4 Rupniu Str., Dzerie, Latvia, LV-2114
Marketed by Unilab Europe
Hamburger S 20251
Hamburg, Germany
Imported and Distributed by
UNILAB, Inc.
Unilab Tower Bldg., Mandaluyong City
Metro Manila, Philippines

Date of Revision: February 2017

Reg.: IP/QP/HL
NO15902706/1