℞

Clomifene citrate

Pregina

50 mg Tablet
Ovulation Inducing Agent

FORMULATION
Each tablet contains:
Clomifene citrate ..... 50 mg

PHARMACOLOGIC CATEGORY
Ovulation-inducing Medicine

PRODUCT DESCRIPTION
Clomifene citrate 50 mg is a white, uncoated tablet.

CLINICAL PHARMACOLOGY
Pharmacodynamics
Clomifene citrate is a nonsteroidal agent which may induce ovulation in appropriately selected anovulatory women. It has both estrogenic and anti-estrogenic properties but its precise mechanism of action in inducing ovulation has not been determined.

Clomifene citrate appears to stimulate the release of the pituitary gonadotropins, follicle-stimulating hormone (FSH) and luteinizing hormone (LH), either from direct stimulation of the hypothalamic-pituitary axis or from a decreased inhibitory influence of estrogens on the hypothalamic-pituitary axis by competitively inhibiting the endogenous estrogens of the uterus, pituitary or hypothalamus. This results in development and maturation of the ovarian follicle, ovulation and subsequent development and function of the corpus luteum.

Clomifene citrate may also directly affect the biosynthesis of ovarian hormones. Clomifene has no progestational, androgenic or antiandrogenic effects and does not appear to interfere with pituitary-adrenal or pituitary-thyroid function.

Pharmacokinetics
Clomifene is readily absorbed from the gastrointestinal tract after oral administration. Blood concentrations of the drugs are at peak enriched isomer concentration in plasma during radiolabeled clomifene citrate. The volume of distribution and plasma protein binding of clomifene citrate is unknown. The drug appears to be less well absorbed and more rapidly eliminated than the 2-isomer.

Clomifene citrate is metabolized in the liver and slowly excreted via the bile. The biological half-life is reported to be 5 days although traces are found in the feces for up to 6 weeks. Mean urinary excretion is apparently 8% and fecal excretion is about 42%.

Limited studies suggest that the drug is excreted slowly from a sequestered enterohepatic recirculation pool or may be stored in body fat and subsequently released slowly.

INDICATION
Clomifene citrate is indicated for induction of ovulation in patients with persistent ovulatory dysfunction who desire pregnancy. The safety and use of treatment of cumulative duration of more than 6 cycles should be supervised by physicians experienced in the management of gynecologic or endocrine disorders. Patients should be carefully selected and monitored with clomifene only after careful diagnostic evaluation. The work-up of the patient must begin with a careful and complete history of menstrual and reproductive function, and a complete physical examination. It should be followed by a selective and careful laboratory investigation based on historical and physical findings.

1. Exclusion of pregnancy: Treatment should be withheld until diagnosis of pregnancy has been ruled out.
2. Assessment of abnormal excessive bleeding: Patients with this condition should be evaluated prior to therapy. It is important to ensure that malignant lesions are not overlooked. Exclusion of presence or history of liver dysfunction: Assess liver function prior to treatment.
3. Exclusion of ovarian cysts: Do not use clomifene citrate in patients with enlarged ovaries entrapment except those with polycystic ovary syndrome (PCOS). Perform pelvic examination before the first and each subsequent treatment cycle.
4. Confirmed ovulatory dysfunction: Establish the diagnosis of ovulatory dysfunction by standard methods, including basal body temperature, cervical mucus, vaginal smears, cervical mucus, endometrial biopsy, and urinary pregnanediol determinations.
5. Exclusion of primary pituitary or ovarian failure: Successful therapy with clomifene requires intact pituitary and ovarian function. Ovulatory dysfunction in the presence of abnormally high levels of pituitary gonadotropins indicates ovarian failure and does not respond to clomifene.
6. Assessment of estrogen levels: Adequate levels of endogenous estrogen provide a favorable prognosis for treatment with clomifene. Physiologic indicators of normal endogenous estrogen include basal body temperature, cervical mucus, endometrial biopsy, assay of urinary estrogens, or from bleeding in response to progesterone. Reduced estrogen level, although less favorable, does not preclude successful therapy.
7. Exclusion of mechanical impediments to conception: Exclude or adequately treat mechanical impediments to conception (e.g., tubal infection) before taking clomifene.
8. Exclusion of medical impediments to pregnancy: Clomifene is not a substitute for appropriate therapy for a coexisting endocrine disorder (e.g., hypothyroidism, adrenal e.g., diabetes, diseases of pituitary origin) that may impede fetal development should be undertaken. Adequately therapy demonstrated only after the underlying disorder has been adequately treated.
9. Exclusion of male factor: Semen analysis should be performed. Abnormal sperm count, motility analysis and other indicated examination. The efficacy of clomifene for male infertility has not been established. Clomifene increases serum and gonadotropins levels in men who are given clomifene citrate. However, the cause and effect relationship between reports of testicular tumors and the administration of clomifene is unknown.

Clomifene has also been used in conjunction with gonadotropins and *in in vitro* fertilization programmes.

Clomifene has also been used in the treatment of male fertility due to oligospermia to stimulate gonadotropin release and enhance spermatogenesis.

DOSAGE AND ADMINISTRATION
General Considerations
The workup and treatment of candidates for clomifene citrate therapy should be supervised by physicians experienced in the management of gynecologic or endocrine disorders. Clomifene should only be given to appropriately chosen anovulatory patients. The therapeutic objective should be balanced with safety to ensure that treatment is discontinued with the patient and others who are involved in the achievement of a pregnancy. The plan of therapy should be outlined in advance.

Data suggest that clomifene treatment of 100 mg/day for 5 days is more effective compared with 50 mg/day for 5 days. However, an increased incidence of ovarian hyperstimulation and other adverse effects may be expected. It is recommended that clomifene citrate therapy start with the lowest effective dose and increased only in those patients who do not respond to the first course of treatment. Special care in dosage is particularly recommended if unusual sensitivity to pituitary gonadotropin is suspected, such as in patients with PCOS.

Recommended Dosage
First Course: 50 mg once a day for five (5) days.
Treatment may be started at any time in the patient who has had no recent uterine bleeding. Start clomifene therapy about the 5th day of the menstrual cycle if progestin-induced bleeding is planned or if spontaneous menstruation occurs prior to therapy. When ovulation does occur at this dose, there is no advantage to increasing the dose in subsequent treatment cycles.

If ovulation appears not to have occurred after the first course of therapy, a second course of 100 mg once a day for five (5) days may be given as a single dose or divided dose for five (5) days. This course may be started as early as 30 days after the previous. **Increasing the dose or duration of therapy beyond 100 mg per day for 5 days should never be undertaken.**

The majority of patients who respond do so during the first course of therapy; three (3) courses should constitute an adequate therapeutic trial. If ovulatory menses have not occurred, the diagnosis should be re-evaluated. Treatment beyond three (3) is not recommended in a patient who does not exhibit evidence of ovulation.

In most patients, ovulation will take place 5 to 12 days after completion of therapy. Coitus should be timed to coincide with the expected time of ovulation. Appropriate tests to determine ovulation may be useful during this period.

Long-term cyclic therapy (more than a total of 6 cycles, including 3 ovulatory cycles) with clomifene is not recommended. The relative safety of long-term cyclic administration of clomifene citrate has not been established.

CONTRAINDICATIONS
- Known hypersensitivity to clomifene citrate or to any ingredient in the product
- Pregnancy
- Active liver disease or history of liver dysfunction
- Uterine abnormal bleeding of undetermined origin
- Ovarian cysts
- Visual disturbances or its association with polycystic ovarian syndrome) and pre-existing endometriosis

WARNINGS AND PRECAUTIONS
Ovarian Hyperstimulation Syndrome (OHSS)
Clomifene citrate alone or in combination with gonadotropins has been reported to cause OHSS. Early warning signs of OHSS include abdominal pain and distention, nausea, vomiting, diarrhea, and weight gain. OHSS may progress from mild to severe with the development of ascites, hydrothorax, hemoconcentration, and hypoproteinemia may occur. In severe cases, OHSS can induce gross ovarian enlargement, thromboembolic phenomena, hypovolemic shock, anuria, and pleural effusion. OHSS has been associated with transient liver function test abnormalities suggestive of hepatic damage. This may be accompanied by morphologically reversible liver lesions.

Severe forms of OHSS have been reported rarely with the use of clomifene (either alone or in combination with gonadotropins). Pericardial effusion, anasarca, hydrothorax, acute abdomen, renal failure, arterial thromboembolic phenomena, intracranial and ovarian hemorrhage, deep venous thrombosis, torsion of the ovary, and acute respiratory distress occurred in these patients. Death due to hypovolemic shock, disseminated intravascular coagulation, and multiorgan failure has occurred. A competition results, rapid progression to the severe form of the syndrome may occur.

The lowest dose consistent with expected clinical results should be used to minimize the hazard associated with ovarian abdominal pain and ovarian enlargement, associated with clomifene citrate therapy. Maximal enlargement of the ovary, whether physiologic or abnormal, may not occur until several days after the last dose of the drug. Some patients with PCOS who are unusually sensitive to gonadotropins may experience an exaggerated response to usual doses of clomifene. Patients with PCOS should be started on the lowest dosage and carefully monitored to determine the duration for the first course of therapy.

Patients should be informed of the possibility of ovarian cyst formation during therapy and should be instructed to return for repeat pelvic examination between 2 to 3 weeks after starting each course of treatment. Patients should be informed that the presence of any abdominal or pelvic pain, weight gain, discomfort or distention after taking clomifene.

Patients who complain of pain in the pelvic or abdominal area should be examined to rule out ovarian cyst formation or other causes. Due to fragility of enlarged ovaries in some cases, abdominal and pelvic examination should be performed carefully.

Unless the ovaries have returned to its pretreatment size, additional dose of clomifene should not be given in patients who develop cysts. Duration of the next course should be shortened. Clomifene-induced ovarian enlargement and cyst formation ordinarily regress spontaneously within a few days or weeks after discontinuation of treatment. Cystic enlargement should always be managed conservatively unless surgical indication exists.

Visual Symptoms
During and/or after visual symptoms such as spots or flashes (scintillating scotomata), diplopia, photophenes, or other visual disturbances may occur occasionally during therapy. These symptoms appear to be related to the dose and duration of treatment and generally disappear within a few days or weeks after discontinuation.

Patients should be warned that visual symptoms may make activities such as driving or operating machinery more hazardous than usual under conditions of variable lighting. These visual symptoms are usually reversible upon cessation of therapy. Cases of prolonged visual disturbance and irreversible especially with increased dosage and duration of therapy. It is recommended to discontinue treatment and a complete ophthalmologic evaluation be performed if a patient has any visual symptoms.

Diagnosis Prior to Clomifene Therapy
The incidence of a twinning disorder and the tendency to develop endometrial carcinoma increases with age. Prior to clomifene therapy in such patients, dilatation and curettage should always be done for diagnosis. Full diagnostic measures are mandatory if abnormal bleeding is present.

Ovarian Cancer
Fertility drugs have been rarely associated with ovarian cancer with infertility itself being a primary risk factor. Since epidemiological data suggest that prolonged therapy with clomifene citrate may increase this risk, it is recommended duration of treatment should not be exceeded.

Uterine Fibroids
Patients with uterine fibroids should use clomifene with caution because of the risk of further enlargement of the fibroids.

Hepatic Impairment
Clomifene is not recommended in these patients (see Contraindications) and therefore clinical evaluation of liver function should always precede therapy.

Pre-existing Mental Depression or Thrombophlebitis
Use clomifene with caution in patients with pre-existing mental depression or thrombophlebitis to avoid this risk of exacerbation of the disease.

Teratogenic/Non-teratogenic Effects: In clinical studies, the overall incidence of birth anomalies resulting from clomifene ingestion during pregnancy was within the range of that reported for the general population.

Pregnancy Wastage: A pregnancy (single and multiple) wastage or fetal loss rate of 21.4% (abortion rate of 15.6% and stillbirth rate of 5.8%) was reported in 1,180 births. Premature rate 0.47%, and pregnancies with one or more still births 1.01% have been observed in patients of all diagnoses during clinical investigation of clomifene citrate.

Multiple Pregnancy: Treatment with clomifene citrate increases the chance of multiple pregnancy (including triplets, quadruplets and quintuplets). The incidence of multiple pregnancy was 7.0% in clinical studies. Among these pregnancies, 6.9% were twin, 0.5% triplet, 0.3% quadruplet, and 0.1% quintuplet. The ratio of maternal age to dosage's twins was 1-9 of the twin pregnancies. Before starting treatment, the patient and her spouse should be advised of the frequency and potential hazards of multiple pregnancy.

Combination therapy with clomifene citrate and chorionic gonadotropin has resulted in simultaneous bilateral tubal pregnancy, an extremely rare form of twin pregnancy.

Ectopic Pregnancy: An increased risk of ectopic pregnancy (including tubal and ovarian sites) has been observed in women who conceive after clomifene therapy.

INTERACTIONS WITH OTHER MEDICAMENTS
None known.

STATEMENT ON USAGE FOR HIGH RISK GROUPS
Pregnancy/ Pregnancy Category X
Clomifene citrate should not be administered during pregnancy. Although no causative evidence of a teratogenic effect for clomifene or the human fetus has been established, birth anomalies (e.g., Down's syndrome, ectrodactyly, club foot, tibial torsion, blocked tear duct, hemangioma, cleft lip and palate) have been reported. These observations in clinical studies have occurred at an incidence within the range reported for the general population.

Before starting treatment, the patient and her spouse should be advised of the frequency and potential hazards of multiple pregnancy.

Lactation: It is not known whether clomifene citrate is excreted in human milk. in some patients, clomifene citrate may suppress lactation.

Infants and Children: Clomifene citrate is not intended for use in children.

Elderly: Clomifene is not intended for use in the elderly.

UNDESIRABLE EFFECTS
The incidence and severity of adverse events due to clomifene citrate therapy are usually related to the dose and duration of use.

Birth Defects: Congenital heart lesions, Down's syndrome, neural tube defects, club foot, congenital gut lesions, hypospadias, microcephaly, hernia and cleft palate, congenital hip, hemangioma, undescended testes, polydactyly, conjoined twins with teratomatous malformation, patent ductus arteriosus, anencephalic monster, arteriovenous fistula, kyphosis, hernia, umbilical hernia, syndactyly, pectus excavatum, myopathy, dermoid cyst of scalp, omphalocele, spina bifida occulta, ichthyosis, congenital heart disease, inguinal hernia, and cardiac defect

Body as a Whole: Fever, tinnitus, weakness, fatigue, increased appetite

Cardiovascular Effects: Arrhythmia, chest pain, edema, hypertension, palpitation, phlebitis, pulmonary embolism, shortness of breath, tachycardia, thrombophlebitis

CNS Effects: Increased nervous tension, insomnia, depression, vertigo, convulsions, dizziness or lightheadedness/vertigo, headache, migraine, paresthesia, stroke, syncope, and seizures; anxiety, irritability, mood changes, and psychosis have also been reported

Dermatologic: Rash, urticaria or allergic dermatitis, acne, allergic reaction, erythema, erythema multiforme, erythema nodosum, hyperhidrosis, reversible hair loss, and alopecia

Gastrointestinal Effects: Abdominal-pelvic discomfort, (distention, bloating, pain, or soreness), nausea and/or vomiting, diarrhea, constipation, and dysgeusia

Genitourinary Effects: Abnormal ovarian enlargement, rare instances of massive ovarian enlargement, ovarian hyperstimulation syndrome, cystic ovarian pain (mittelschmerz), and cyst formation (usually reversible). Rare cases of endometrioma and exacerbation of pre-existing endometriosis have also been reported

Multiple pregnancies (but rarely more than twins), including simultaneous intrauterine and extrauterine pregnancies, hydatidiform mole, hydatid pregnancy, ectopic or interstitial pregnancies, and uterine hemorrhage have also been reported.

Hepatic Effects: Abnormal liver function tests, jaundice, hepatitis, and infectious hepatitis

Musculoskeletal Effects: Arthralgia, back pain, myalgia

Special Senses: Dose-related visual symptoms described as "blurring" or spots or flashes (scintillating scotomata) usually appear due to intensification and prolongation of after-images, ophthalmoplegia, photophobia, photopsia, photophenes, diplopia, phosphenes, and reduced visual acuity. Cataracts and optic neuritis have been reported rarely. Light-sensitivity, light floaters, tears, etc., and other ocular symptoms not otherwise specified have also been reported.

Tumors/Neoplasms: Ovarian cancer, liver cancer hepatic hemangiosarcoma, liver cell adenoma, hepatocellular carcinoma, breast (ductal)cystic disease, breast carcinoma), endometrium endometrial carcinoma, nervous system (astrocytoma, pituitary tumor, protactinoma, neurofibromatosis, glioblastoma multiforme, brain abscess), trophoblastic (trophoblastic mole, choriocarcinoma), miscellaneous (hepatoma, pituitary tumor, renal cell cancer, renal carcinoma, Hodgkin's lymphoma, tongue carcinoma, bladder carcinoma), and neoplasms of offspring (neuroectodermal tumor, thyroid tumor, hepatoblastoma, nephroblastoma, leukemia)

Isolated reports of endocrine-related or dependent tumors/neoplasms or their aggravation

Vasomotor Symptoms: Hot flushes resembling menopausal symptoms, breast discomfort, breast soreness, breast tenderness, breast tumor menses/menorrhagia, intermenstrual spotting, vaginal dryness, and decrease in libido

Other Effects: Leukocytosis increased, urinary frequency or volume, weight gain, weight loss, edema, moderate and reversible hair loss, prolonged continuous therapy and thyroid disorder

Laboratory Studies: Administration of clomifene resulted in retention of sulfobromophthalein (BSP) in patients in whom BSP is measured. Retention was usually minimal unless associated with prolonged continuous use. Clomifene is not apparently correlated liver disease.

Clomifene has not been shown to cause significant abnormality in hematologic or renal tests, in protein bound iodine or in serum cholesterol levels.

OVERDOSAGE & MANAGEMENT
There is no experience with overdosage. There is no known antidote. Overdosage of clomifene citrate may be reported to the FDA at www.fda.gov.ph AND Unilab at (+632) 858-1000 or productsafetyfdunilab.com.ph. By reporting undesirable effects, you can help provide more information on the safety of this medicine.

STORAGE CONDITIONS
Store at temperatures below 30°C.
Protect from heat, light, and excess humidity.
Keep the product out of reach and sight of children.
Store in original packaging.

CAUTION
Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

AVAILABILITY
PVC blister pack of 10's; Box of 10 tablets.

Manufactured by:
RENATA LIMITED
Section VB Mile Ala Road
Mirpur, Dhaka, Bangladesh
Imported and Distributed by:
UNILAB, Inc.
Unilab Tower, 191 United Street, Mandaluyong City
Metro Manila, Philippines

Reg. IPOPHL

BF156837R03