Hepatitis-B Vaccine (recombinant DNA)

Revac-B®
20 mcg /mL
Suspension for Injection (IM)

FORMULATION
Each mL single-dose (Adult Dose) vial contains:
Active: Hepatitis B Surface Antigen ..... 20 mcg
Aluminum Hydroxide Gel (as Al^{+3}) ..... 0.5 mg
Thiomersal (as preservative) ..... 0.05 mg

PRODUCT DESCRIPTION
Recombinant Hepatitis B vaccine (Revac-B®) is a sterile suspension for injection containing purified, non-infectious major surface antigen of the hepatitis B virus (HBsAg) produced by means of recombinant DNA technology. The antigen is adsorbed onto high affinity aluminum hydroxide gel medium. Hence, the suspension appears almost white and translucent. The vaccine fulfills the requirements of the World Health Assembly for recombinant DNA vaccines.

Recombinant Hepatitis B vaccine (Revac-B®) is presented in 0.5ml * 1 glass vial. The vaccine upon storage may present a fine white deposit with a clear colorless supernatant. The presence is slightly opaque once shaken.

Recombinant Technology
The Hepatitis B surface antigen (HBsAg) is produced by yeast cells (Hansenula sp.) in which the HBsAg particles which carry the gene that codes for the major surface antigen protein of the HBV. The HBsAg expressed in yeast cells is purified by complex physical, chemical and biochemical processes. The HBsAg particles thus produced are highly immunogenic non-infectious HBV-like spherical particles of an average diameter of 20-24 nm containing non-glycosylated polypeptides in a lipid matrix. **No addition of human origin is used in the manufacture of HBsAg protein.**

CLINICAL PHARMACOLOGY
Hepatitis B virus (HBV) is a DNA virus consisting of antigenically distinct surface and core components. The major antigenic determinant of the viral surface is HBsAg. The presence of HBsAg in serum indicates the presence of an acute or chronic HBV infection or a chronic carrier state.

Hepatitis B virus is one of several hepatitis viruses that can cause inflammation of the liver, which in some pathologic changes in the liver. The incubation period for HBV infection after exposure to the virus to onset of symptoms is usually 6 to 8 months.

Hepatitis B virus infection can have serious consequences including acute massive hepatic necrosis, cirrhosis of the liver, chronic liver disease and even hepatocellular carcinoma (HCC). Studies have shown a clear link between chronic hepatitis infection and development of cirrhosis and 80% of HCCs are caused by HBV. Hepatitis B vaccine has been shown to reduce the incidence of hepatocellular carcinoma because it can prevent many liver cancers.

Hepatitis B vaccine stimulates active immunity to HBV infection. The HBsAg, which is present in vaccine, induces the production of specific antibody (production of antibody to HBsAg (anti-HBs)). anti-HBs neutralize HBV so that its pathogenic or infective properties are inhibited.

Protection against HBV infection is virtually complete in individuals who develop adequate antibody levels after hepatitis B vaccine administration. The level of antibody considered as need for booster dose after primary immunization with hepatitis B vaccine has not been fully determined.

Immunogenicity in Adults:
Results of a study in adults (n=226), with a mean age of 27.65 years (M, given a 3-dose regimen (0, 30th and 60th days) of 20 mcg doses of a recombinant hepatitis B vaccine derived from *Pichia pastoris* yeast showed seroconversion rates of 100% by day 60 and 100% by the 90th day of the study. The seroprotection rates were 71% by the 30th day, 99.6% by the 60th day, and 100% by the 90th day. The geometric mean antibody titer (GMT) increased from 1,000 mIU/mL on day 30 to 3,000 mIU/mL on day 60 and remained at that level for 6 months as per data recorded. These results were comparable with two other commercially available recombinant hepatitis B vaccines (one derived from *Saccharomyces cerevisiae* yeast and the other from plasma).

Immunogenicity in Patients with Hepatitis C virus (HCV) Infection:
The immunologic response to hepatitis B vaccine may be decreased in patients with HCV infection and may be adversely affected by HCV viral overload.

Immunogenicity in Hemodialysis and Immunocompromised Patients
In general, the immunologic response to hepatitis B vaccine in chronic hemodialysis patients and other immunocompromised individuals, including adults, and adequate levels of protection, persist for shorter periods.

INDICATION
Hepatitis B vaccine 20 mcg/mL dose is for the active immunization against hepatitis B infection in adults and children >15 years old.

- For primary immunization against HBV infection caused by all known subtypes, and to stimulate active immunity to HBV in individuals, who are, or will be, at increased risk of potential exposure to HBV or HBsAg-positive materials (e.g., blood, plasma, serum).

- Hepatitis D occurs only in the presence of hepatitis B. By immunizing against hepatitis B, hepatitis D infection may be prevented.

- Hepatitis B vaccine will not prevent hepatitis caused by other agents, such as hepatitis A, C, E, or unidentified pathogens known to infect the liver.

DOSAGE AND ADMINISTRATION
For deep intramuscular (IM) injection into the deltoid muscle in adults and adolescents.

- **Do not inject intravenously or intradermally.** The vaccine may be administered subcutaneously (SC) in patients with thrombocytopenia or coagulation disorders.

- Do not inject into the gluteal muscle since this route of administration may result in lower immune response.

A. Primary Immunization Schedule
1st dose: On selected date one
2nd dose: 30 days after the first dose
3rd dose: 180 days after the first dose

Note: This vaccine may be used to complete a vaccination course in persons who have received previously received another hepatitis B vaccine. Booster dose is not routinely recommended.

B. Special Dosage Recommendations
- Post-exposure prophylaxis: Persons accidental to HBV infection but still negative for HBsAg and IgM anti-BC.

1st dose:	On selected date
2nd dose:	30 days after the first dose
3rd dose	60 days after the first dose
4th dose:	1 year after the first dose

- Immunocompromised patients will require additional dosage as follows:

1st dose of 40 mcg (2ml): On selected date
2nd dose of 40 mcg (2ml): 30 days after the first dose
3rd dose of 40 mcg (2ml): 60 days after the first dose
4th dose of 40 mcg (2ml): 180 days after the first dose

Or, as prescribed by a physician.

Directions for Use
- Inspect visually for particulate matter and discoloration prior to administration.
- The vaccine should be well shaken before use to obtain a uniform, turbid suspension.
- No dilution or reconstitution is necessary.
- Recombinant Hepatitis B vaccine (Revac-B®) should not be mixed with other vaccines.
- Strict aseptic technique should be observed when drawing up the contents of the vial. If contaminated, it has the potential to become a source of infection to patients.
- Any vaccine remaining in a single-dose vial should be discarded properly.

Stability and Potency:
- Recombinant Hepatitis B vaccine (Revac-B®) is stable and potent for 36 months at +2°C to +8°C. Brief temperature excursions may be permissible at 37°C for 1 month and 45°C for 1 week did not result in the loss of its immunogenicity.

CONTRAINDICATIONS
- Known hypersensitivity to hepatitis B-containing vaccine or any component of the vaccine, including yeast.
- Severe febrile illness.

WARNINGS AND PRECAUTIONS
- Prior to immunization, the patient's medical history should be reviewed for any previous sensitivity to the vaccine. As with any vaccine, appropriate medical treatment including epinephrine should be available for immediate use in case of an anaphylactoid reaction.
- Patients who develop hypersensitivity reactions after an injection should not receive further injection of the vaccine.
- Syncopal reactions may occur since thiomersal has been used as the preservative in the process of this vaccine and residues of it are present in the final product.
- Any serious active infection including febrile illness is a reason for postponing use of the vaccine except when, in the opinion of the physician, withholding the vaccine entails a greater risk to the patient.
- Use with caution in individuals with severely compromised cardiopulmonary status or to those who may not tolerate a transient systemic reaction should pose a significant risk.
- Exacerbation of multiple sclerosis has been rarely reported after hepatitis B vaccine administration; no causal relationship has been established. Thus, in patients with multiple sclerosis, the benefit of immunization for prevention of hepatitis B infection and the potential risk should be carefully considered before the risk of exacerbation of the disease.
- Immunocompromised and hemodialysis patients may require a booster dose to maintain their healthy individuals to indicate adequate circulating antibody titers (**see Special Dosage Recommendations**). For those receiving immunosuppressive therapy, deferral of vaccination until the therapy is completed may be considered (**see Interactions with other Other Medicaments**).
- Since hepatitis B virus has a long incubation period to manifest the symptoms of infection, some patients may exist in the vaccine while the infection remains unrecognized. In such cases, the vaccine may not prevent HBV infection. Moreover, it may not prevent infection in individuals who do not achieve protective antibody titers.
- A number of factors have been associated with the reduction in immune response to hepatitis B vaccines. These include older age, male gender, obesity, smoking, use of immunosuppression and some chronic underlying diseases. Serological testing after the vaccination series in individuals who may be at risk of not achieving seroprotection after a complete course of hepatitis B vaccine.
- Patients with chronic liver disease or with HIV infection or hepatitis C carrier should not be precluded from vaccination against hepatitis B. Such patients may be at risk for acquiring additional infection and the vaccine is therefore recommended in a case to case basis by the physician.
- As with any vaccine, a protective immune response may not be obtained in all vaccines.

INTERACTIONS WITH OTHER MEDICAMENTS
- **Hepatitis B Immunoglobulin (HBig)** To avoid interference with the active immune response, it should not be administered in the same syringe and should not be injected at the same site.
- **Other Vaccines** (e.g., vaccine against Hepatitis A, Diphtheria, Tetanus, Pertussis, Influenza, Polio, BCG, polio, measles, mumps, rubella, diphtheria, tetanus, pertussis, influenza, pneumococcal): In general, simultaneous administration of these vaccines has not been found to interfere with the immune response. An increased adverse effect of vaccine uses and syringes should be used for simultaneous administration of injectable vaccines.
- **Anti-infective agents**: In general, concomitant therapy with these agents will not affect the immune response to inactivated vaccines, including hepatitis B vaccine.
- **Immunosuppressive agents** (e.g., corticosteroids, cytotoxic alkylating agents, antimetabolites, radiation) individuals receiving such agents may have a reduced response.

STATEMENT ON USAGE FOR HIGH RISK GROUPS
- **Pregnancy/ Pregnancy Category C:** There are no well-controlled studies in pregnant women. Hepatitis B vaccine should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.
- **Lactation:** It is not known whether hepatitis B vaccine is excreted in human milk. Since many medicines are excreted in human milk, the vaccine should be used with caution in breastfeeding women.
- **Children:** Hepatitis B vaccine is well-tolerated and highly immunogenic in children.
- **Elderly:** Hepatitis B vaccine is less immunogenic in adults >60 years of age when compared with younger patients. No overall differences in safety were observed between the elderly and younger adult subjects.

UNFAVORABLE EFFECTS
Recombinant Hepatitis B vaccine is generally well tolerated. The following undesirable effects have been reported after the use of Hepatitis B vaccine; however, no causal relationship has been established to the vaccine has not been established:

Injection Site Reactions: Burning, erythroderm, erythema, induration, inflammation/swelling, focal pain, nodule formation, pruritus, soreness, tenderness, warmth.

Body as a Whole: Achiness, chills, fatigue/asthenia, fever (≥37.8°C), flushing, influenza-like symptoms, malaise, paresis, tingling, sweating, sensation of warmth, weakness.

Dermatologic/Hypersensitivity Reactions: Anaphylaxis-like symptoms of hypersensitivity reactions including rash, pruritus, urticaria, angioedema, bronchospasm, dyspnea, bronchitis, bronchial disorder, pharyngitis/throat symptoms), papulation or symptoms consistent with a hypersensitivity episode have been reported within the first few hours after administration of hepatitis B vaccine. Erythema multiforme, Stevens-Johnson syndrome, petechiae, and thrombocytopenic reaction with delayed onset has been reported days to weeks after administration of the vaccine. This reaction consists of dermatologic reactions such as erythema multiforme, Stevens-Johnson syndrome, syndrome) and cutaneous vasculitis purpura, polyarteritis nodosa, systemic lupus erythematosus (SLE), baboche erythema.

Musculoskeletal System: Arthralgia (including temporomandibular), arthritis, myalgia, pain in arm, shoulder, back and/or neck, myalgia, muscle weakness

Cardiovascular System: Hypotension, syncope, palpitations, tachycardia

Nervous System: Agitation, convulsions, demyelinating disease of the CNS, disturbed sleep/insomnia, encephalitis, encephalopathy, febrile convulsions, hallucinations, headache, hyperkinesthesias, meningitis, neuritis, multiple sclerosis, exacerbation of multiple sclerosis, neuritis including Guillain-Barré syndrome, optic neuritis, myelitis including transverse myelitis), paresthesia, peripheral neuritis, Bell's palsy, somnolence, vertigo

Gastrointestinal System: Abdominal pain/cramps, constipation, diarrhea, dyspepsia, nausea, nausea, vomiting

Hematologic: Increased erythrocyte sedimentation rate, lymphadenopathy, thrombocytopenia

Immune System: Anaphylaxis, chills, fatigue/asthenia, fever, serum sickness-like reactions, urticaria

Respiratory System: Rhinitis/coryza (including asthma-like symptoms), cough, influenza, moaldeas, pharyngitis, rhinitis, symptoms of upper respiratory disease

Gastrointestinal Disorders: Abdominal pain/cramps, constipation, diarrhea, diminished appetite/anorexia, dyspepsia, nausea, vomiting

Urogenital System: Dysuria

Hepatobiliary Disorders: Abnormal liver function tests

Hematologic: Increased erythrocyte sedimentation rate, lymphadenopathy, thrombocytopenia

Special Senses: Earache, tinnitus, conjunctivitis, keratitis, optic neuritis, uveal disturbances

OVERDOSE AND TREATMENT
No case of overdose has been reported.

STORAGE CONDITIONS
- Store at 2°C to 8°C.
- Do not freeze. Discard if product has been frozen.
- Keep the product out of sight and reach of children.

AVAILABILITY
Box of 1's (1 mL vial)

CAUTION: Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

DATE OF REVISION OF THE PACKAGE INSERT
May 2018

Manufactured by
Bharat Biotech International Ltd.
Shamirpet Mandal
Ranga Reddy District
Hyderabad-India

Imported and Distributed by
United Laboratories Inc.
Km. 18 West Service Rd., Muntinlupa City
Philippines

Under Authority by
ULVaccines Inc.
Km. 18 West Service Rd.
Muntinlupa City
Philippines

ULVaccines

Revac-B-Insert