GLICLAZIDE

Gliclazide Glubitor® 80 mg Tablet
oral hypoglycemic

Rx

FORMULATION
Each tablet contains 80 mg of gliclazide.

PHARMACEUTICAL FORM
White, round, flat, bevelled-edged tablet, with a double score on one side and "S/P" imprint on the other side.

CLINICAL PARTICULARS
Therapeutic indications
For insulin-dependent diabetes (type 2) in adults when dietary measures, physical exercise and weight loss alone are not sufficient to control blood glucose.

DOSAGE AND ADMINISTRATION
Dosage
The dose may vary from 40 mg to 320 mg (1 to 4 tablets per day, or from 30 to 120 mg taken orally in a single intake at breakfast time.
Tablets should be taken whole with a glass of fluid.
If you have forgotten to take a medicine at the time to take the dose next day, as with any hypoglycemic agent, the dose should be adjusted according to the individual blood glucose monitoring results and HbA1c level.

Initial dose
The recommended starting dose is 30 mg daily.

If blood glucose is effectively controlled, this dose may be used for maintenance treatment. If blood glucose is not adequately controlled, the dose may be increased to 60 mg, 90 mg or 120 mg daily. The dose should be increased in steps of one tablet every month, **except if month's level of glycosylated hemoglobin** has not decreased by at least after two weeks of treatment. In such cases, the dose may be increased at the end of the second week of treatment.

The maximum recommended daily dose is 120 mg.

Switching from tablets containing 80 mg of gliclazide to gliclazide modified-release tablets
1 tablet containing 80 mg of gliclazide is comparable to 1 modified-release tablet of gliclazide 30 mg. Switching from one to the other should be carried out without delay.

Switching from another oral antidiabetic agent to gliclazide
Gliclazide can be used to replace other oral antidiabetic agents. The dosage and the half-life of the previous antidiabetic agent should be taken into consideration when switching to gliclazide.

A transitional period is not generally necessary. A starting dose of 30 mg should be used and this should be adjusted to suit the patient's blood glucose response, as described above.

When switching from a hypoglycemic sulfonylurea with prolonged half-life, a washout period of several days may be necessary to avoid additive effects which may result in hypoglycemia. The initial dose of gliclazide should be as mentioned above and titration should be carried out as mentioned above, according to the patient's blood glucose response.

Switching from another oral antidiabetic agent to gliclazide
Gliclazide can be used in combination with biguanides, alpha glucosidase inhibitors and insulin. In patients not adequately controlled on gliclazide monotherapy, the dose of gliclazide may be titrated under close medical supervision.

Special populations
Elderly people
Gliclazide should be prescribed using the same dosing regimen recommended for patients under 65 years of age.

Patients with renal impairment
In patients with mild to moderate renal impairment, the same dosing regimen can be used as in patients with normal renal function with careful patient monitoring. These data have been confirmed in clinical trials.

Patients at risk of hypoglycemia
Undernourished or malnourished
Severe or poorly compensated endocrine disorders (hypothalamic, hypothyroidism, adrenocorticotropic deficiency)
Withdrawal of prolonged and/or high dose corticosteroid therapy
Severe vascular disease (severe coronary heart disease, severe carotid impairment, diffuse vascular disease).

It is recommended that the minimum starting dose of 30 mg is used.

Pediatric population
The safety and efficacy of gliclazide in children and adolescents have not been established. No data are available.

CONTRAINDICATIONS
This medicinal product is contraindicated in case of:
• hypersensitivity to gliclazide or to any of the excipients, other sulfonylureas, or sulfonamides,
• type 1 diabetes,
• diabetic pre-coma and coma, diabetic ketoacidosis,
• severe renal or hepatic failure (in these cases the use of insulin is recommended),
• lactation (as this medicinal product **INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION**).
• lactation (see **PREGNANCY AND LACTATION**).

SPECIAL WARNINGS AND PRECAUTIONS FOR USE
Hypoglycemia
Gliclazide should be prescribed with caution if the patient is likely to have a regular food intake (including breakfast). It is important to have a regular carbohydrate intake due to the risk of hypoglycemia. Glucose must be taken immediately. A sugar supplement must be fed to consumed or the food is low in carbohydrates.

Hypoglycemia is more likely to occur when the medicine starts, following prolonged or strenuous exercise, alcohol intake or if a combination of hypoglycemic agents is being taken.

Hypoglycemia may occur following administration of sulfonylureas (see **UNDESIRABLE EFFECTS**). These reactions may be severe and prolonged. Hospitalization may be necessary and glucose administration may need to be continued for several days.

Careful monitoring of the patient's blood glucose level, and other patient directions are necessary to reduce the risk of hypoglycemic episodes.

Factors which increase the risk of hypoglycemia
• Patient refusal or inability (elderly patients) to be unable to cooperate
• Malnutrition, irregular schedules, skipping meals or fasting, changes in dietary changes
• Imbalance between physical exercise and carbohydrate intake
• Renal impairment
• Severe hepatic impairment
• Overdose of gliclazide
• Certain endocrine disorders thyroid disorders, hypopituitarism and adrenal insufficiency
• Concomitant administration of certain other medicines (see **INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION**).

Renal and hepatic insufficiency
The pharmacokinetic and pharmacodynamics of gliclazide may be altered in patients with hepatic insufficiency or severe renal failure. A hypoglycemic episode occurring in these patients may be prolonged, so appropriate management should be instituted.

Loss of information
The risks of hypoglycemia, together with its symptoms (see **UNDESIRABLE EFFECTS**), may occur in patients who do not respond adequately, who are irregular in attendance for patient control or who are receiving a low dose.

The patient should be informed of the importance of following dietary advice, of taking regular exercise, and of regular monitoring of blood glucose levels.

Poor blood glucose control
Blood glucose control in a patient receiving antidiabetic treatment may be affected by any of the following: fever, trauma, infection or surgery (see **INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION**). In these cases, short-term insulin therapy is recommended.

The hypoglycemic efficacy of any oral antidiabetic agent, including gliclazide, is attenuated over time in many patients; this may be due to progression in the severity of the disease or to a decreased response to the medicine. This phenomenon is known as secondary failure to distinguish it from primary failure, in which the medicine is ineffective as first-line treatment. Adequate dose adjustment and dietary compliance should be monitored carefully to minimize the risk of secondary failure.

Cardiovascular risk
Concomitant administration of certain other medicines (see **INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION**).

Cardiogenetics
Sulfonylureas in blood glucose, including hypoglycemia and hyperglycemia have been reported, in diabetic patients receiving concomitant treatment with fluoroquinolones, especially in elderly patients. Caution should be exercised when using these combinations.

Renal and hepatic insufficiency
The pharmacokinetic and pharmacodynamics of gliclazide may be altered in patients with hepatic insufficiency or severe renal failure. A hypoglycemic episode occurring in these patients may be prolonged, so appropriate management should be instituted.

Laboratory tests
Determination of glycated hemoglobin levels (or fasting venous plasma glucose) is recommended in assessing blood glucose control. Blood glucose self-monitoring may be helpful.

Treatment of patients with G6PD-deficiency with sulfonylurea agents can lead to hemolytic anemia. Because gliclazide belongs to the class of sulfonylurea agents, caution should be used in patients with G6PD-deficiency and a nonsulfonylurea alternative should be considered.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION
The following medicines may increase the hypoglycemic effect of gliclazide:
Contraindicated combination
• **Miconazole (systemic route, oromucosal gel):** increased risk of hypoglycemic effect with possible onset of hypoglycemic symptoms or even coma.

Combination not recommended
• **Phenylbutazone (systemic route):** increases the hypoglycemic effect of sulfonylureas (displaces their binding from plasma proteins) and also reduces the clearance of the sulfonylurea. It is preferable to use a different anti-inflammatory agent, or else to warn the patient and to reinforce monitoring of blood glucose levels.
• **Fluconazole:** increased risk of hypoglycemia. Blood glucose monitoring must be reinforced.
• **Alcohol:** increases the hypoglycemic reaction (by inhibiting compensatory reactions) that can lead to the onset of hypoglycemic coma. Alcohol or medicines containing alcohol must be avoided.

Combinations requiring precautions for use
• **Beta-blockers, clonidine, guanfacine:** beta-blocking effect and thus, in some instances, hypoglycemia. Monitoring is reinforced by the dose of the antidiabetic agent during and after treatment with danazol.

Combinations taking into consideration during use
• **Chloramphenicol, coumarin derivatives, aspirin, NSAIDs (>150 mg per day of acetylsalicylic acid), sulfonamides, MAOIs, fenfluramine, ACE inhibitors:** there is a risk of a potent and emphasize the importance of blood glucose monitoring. It may be necessary to adjust the dose of the hypoglycemic active substance during and after treatment with the non-steroidal agent.

• **Glucocorticoids (systemic and local route: intra-articular, cutaneous and rectal preparations) and tetracosactide:** increases in blood glucose levels with possible ketosis (reduced tolerance to carbohydrates due to glucocorticoids). Blood glucose monitoring is reinforced, especially at the beginning of treatment. Dose adjustment of the antidiabetic agent may be necessary during and after treatment. If this is necessary to adjust the dose of the antidiabetic active substance during and after treatment with glucocorticoids.

• **Ritodrine, salbutamol, terbutaline (IV):** increased blood glucose levels due to the beta-2 stimulating action. Blood glucose monitoring is reinforced, especially at the start. If necessary, switch to insulin.

Can affect the hypoglycemic control preparations: Gliclazide exposure is decreased by St. John's Wort (*Hypericum perforatum*). Emphasize the importance of blood glucose level monitoring.

The following products may cause dyslycemia
• **Thiazides and other diuretics:** risk of hyperglycemia.
• **Fluoroquinolones:** In case of a concomitant use of gliclazide and a fluoroquinolone, the patient should be warned of the risk of hyperglycemia, and the importance of blood glucose monitoring should be emphasized.

Concomitant use with danazol should be avoided.

PREGNANCY AND LACTATION
Pregnancy
Due to the limited amount of data these 300 pregnancy outcomes from the use of gliclazide during pregnancy, and the risk of hypoglycemia in the fetus and neonate.

Studies in animals have shown reproductive toxicity (see **PRECLINICAL SAFETY DATA**).

As a precautionary measure, it is preferable to avoid the use of gliclazide during pregnancy.

Control of diabetes should be obtained before the time of conception to reduce the risk of congenital abnormalities associated with poor control.

Oral hypoglycemic agents are not suitable; insulin is the drug of first choice for treatment of diabetes during pregnancy. If the patient wishes to become pregnant, therapy is changed to insulin before pregnancy is attempted, or as soon as pregnancy is discovered.

Breastfeeding
It is unknown whether gliclazide/metabolites are excreted in human milk. Given the risk of neonatal hypoglycemia, the product is therefore contraindicated in breastfeeding mothers. Due to its half-life, switching to insulin is recommended.

Fertility
There is no fertility or reproductive performance was noted in male and female rats (see **PRECLINICAL SAFETY DATA**).

Effects on ability to drive and use machines
Gliclazide has no or negligible influence on the ability to drive and use machines. However, patients should be aware of the risk of hypoglycemia or hyperglycemia and its influence on the ability to drive. Caution should be exercised at the beginning of treatment (see **SPECIAL WARNINGS AND PRECAUTIONS FOR USE**).

UNDESIRABLE EFFECTS
The following adverse drug reactions/effects the following undesirable effects have been reported:

Blood and lymphatic system disorders
Very rare: decreased hemoglobin, decreased white blood cell count, decreased platelets. If necessary, switch to insulin.

Immune system disorders
Very rare: allergic reactions (vasculitis, skin rash, photosensitivity, anaphylactic shock).

Metabolism and nutrition disorders
Very rare: hypoglycemia (see **SPECIAL WARNINGS AND PRECAUTIONS FOR USE**).

Nervous system disorders
Very rare: dizziness, tremor, drowsiness, asthenia, headache, paresthesia, convulsions, impaired consciousness, possibly coma and lethal outcome.

Vascular disorders
Very rare: vasculitis.

Gastrointestinal disorders
Very rare: abdominal pain, nausea, vomiting, diarrhea, constipation.

Hepatobiliary disorders
Very rare: hepatic enzymes (AST, ALT, alkaline phosphatase), hepatitis (isolated reports). Discontinue treatment if cholestatic jaundice appears.

Skin and subcutaneous tissue disorders
Very rare: pruritus, urticaria, angioedema, erythema, maculopapular rashes, bullous reactions (such as Stevens-Johnson syndrome, toxic epidermal necrolysis), and exceptionally Drug Rash with Eosinophilia and Systemic Symptoms (DRESS).

Musculoskeletal and connective tissue disorders
Very rare: myalgia.

Eye disorders
Very rare: transient disturbances may occur especially on initiation of treatment, due to changes in blood glucose levels.

Class attribution effects
As for other sulfonylureas, the following adverse events have been observed: cases of hypoglycemia with loss of consciousness, coma, seizures, shallow breathing, bradycardia, hypotension, diaphoresis, skin pallor, headache, nausea, vomiting, drowsiness, loss of rich, cholestasis and jaundice) and hepatitis which regressed after withdrawal of the medicine, and severe liver failure (cholestasis and jaundice).

OVERDOSE
An overdose of sulfonylureas may cause hypoglycemia.

Moderate symptoms of hypoglycemia, without any loss of consciousness or neurological signs, must be corrected by carbohydrate intake, dose adjustment and/or fasting. These symptoms usually disappear after the consumption of sugar, but the patient must continue to be monitored.

Severe hypoglycemic reactions, with coma, convulsions or other neurological disorders are possible and must be treated as a medical emergency, requiring immediate hospitalization.

If hypoglycemic coma is diagnosed or suspected, the patient should be given a rapid IV injection of concentrated glucose solution (50 ml of 10% dextrose). This should be followed by a continuous dextrose infusion at a lower concentration (5%) at a rate that will maintain blood glucose levels above 1g/l. The patients should be monitored closely and, depending on the patient's condition, close medical follow-up may be advised for 24 hours or more.

Gliclazide is not dialyzable by hemodialysis.

PHARMACOLOGICAL PROPERTIES
Pharmacodynamic properties
Pharmacotherapeutic group: Drugs used in diabetes, blood glucose lowering drugs, oral hypoglycemic agents.
ATC code: A10BB09

Gliclazide is hypoglycemic sulfonylurea with antidiabetic active substance differing from other sulfonylureas by having a chemical structure containing both a methyl and a heterocyclic group.

Gliclazide reduces blood glucose levels by stimulating insulin secretion from the B-cells of the islets of Langerhans. It also improves the early phase of insulin secretion in response to glucose.

Gliclazide has a beneficial effect on vascular disease. Gliclazide has hemovascular properties.

Effects on insulin release
In healthy subjects, gliclazide restores the first peak of insulin secretion in response to glucose and increases the second phase of insulin secretion. A significant increase in stimulated insulin secretion is seen 2 hours after administration of a single dose.

Hemovascular properties
Gliclazide also normalizes by two mechanisms, which may be involved in complications of diabetes:

• a partial inhibition of **platelet** aggregation and adhesion, with a decrease in the markers of platelet activation (beta-thromboglobulin, thromboxane B2),
• an action on the vascular endothelium fibrinolytic activity with an increase in tPA activity.

Pharmacokinetic properties
Absorption
Gliclazide is completely absorbed following the dose 6 hours, reaching a plateau, which is maintained from the sixth to the twelfth hour after administration. Intra-individual variability in absorption is low. Absorption is completely absorbed. Food intake does not affect the rate or degree of absorption.

Distribution
Plasma protein binding is approximately 95%. The volume of distribution is around 30 liters. In vitro data makes maintains effective gliclazide plasma concentrations over 24 hours.

Biotransformation
Gliclazide is nearly metabolized in the liver and excreted in the urine less than 1% of the unchanged form is found in the urine. No active metabolites have been detected in plasma.

Elimination
The elimination half-life of gliclazide varies between 12 and 20 hours.

Linearity/Non-linearity
The relationship between the dose administered ranging up to 120 mg and the area under the concentration/time curve is linear.

Special populations
Elderly people
No clinically significant changes in pharmacokinetic parameters have been observed in elderly patients.

Preclinical safety data
Acute toxicity studies in special hazards for humans based on conventional studies of repeated dose toxicity and genotoxicity. Long term carcinogenicity studies have not been done. No teratogenic effects were observed in animal studies. Rats fed with fetuses body weight was observed in animals receiving doses 25 fold higher than the maximum recommended dose in humans.

PHARMACEUTICAL PARTICULARS
Incompatibilities
Not applicable.

STORE AT TEMPERATURES NOT EXCEEDING 30°C
DRUGS SHOULD BE KEPT OUT OF THE REACH OF CHILDREN

CAUTION
Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.

For suspected adverse drug reaction, send report to **UNILAB, Inc.** Medical Information Unit, Tel. No. (632) 8-8888888 ext. 8888888 or (632) 8-8888888 or Fax No. (632) 8-8888888 or 1-800-10-UNILAB (8-645-2522) or e-mail pvcu@unilab.ph.com.ph.
By reporting undesirable effects, you can help provide more information on the safety of the medicine.

AVAILABILITY
Gliclazide (Glubitor®) 80 mg Tablet, in blister pack x 10s (box of 6's).

Manufactured by Interphil Laboratories, Inc.
Km. 23 West Service Road, South Superhighway, Cupang, Muntinlupa City
For Sandoz Philippine Corporation
5th and 6th Floor Ayala North Exchange Tower 1 PKG, Ayala Ave.
cor. Salcedo and Amorsolo, Sts., Bgy. San Lorenzo, Makati, Metro Manila
Distributed by: **UNILAB, Inc.**
Mfg. Licensing and Marketing Group
Metro Manila Philippines

Date of Revision: February 2020
CDS0

Date of First Authorization: September 2012
DRP-1256-03

Reg. CDRR
LI154602R03