AZITHROMYCIN MONOHYDRATE

ZENITH® 500 mg Film-coated Tablet

AMTAC/TEVA/CEBU

FORMULATION
Each film-coated tablet contains:
Azithromycin monohydrate ..... 500 mg

PHARMACEUTICAL FORM
Film-coated tablet, white, debossed with 'ZEN' above the line on one side, scored on the other side.

CLINICAL PARTICULARS
Therapeutic indications
Zenith is indicated for the following infections caused by microorganisms sensitive to azithromycin: Upper respiratory tract infections such as pharyngitis/tonsillitis and sinusitis (as monotherapy).
Lower respiratory tract infections such as acute bacterial exacerbations of chronic obstructive pulmonary disease (comparable control studies) or *Chlamydophila pneumoniae* pneumonia.
Acute exacerbation of bronchitis.
Mild to moderate infections of skin and soft tissue such as erysipelas, impetigo.
Genital infections such as cervicitis or urethritis due to *Chlamydia trachomatis*.
Initial treatment of pelvic inflammatory disease may be combined with other appropriate antibiotics.
Ulcerative genital infections due to *Haemophilus ducreyi* (chancroid).
Zenith is also indicated for the treatment of early Lyme disease (erythema migrans) due to *Borrelia burgdorferi*.

Posology
Recommended dosage schedule due to *Chlamydia trachomatis* or sensitive *Neisseria gonorrhoeae* infections: 1 g as a single dose. Alternatively, 500 mg on day 1, then 250 mg once daily for 4 days.

Other infections
Recommended dosage schedule: 500 mg as a single dose once daily for 3 days. The tablets may be taken with or without food.

Alternatively, for mild to moderate infections, a single dose of 1.5 g (3 tablets) may be given.

Contraindications
Hypersensitivity to the active substance, erythromycin, any macrolide or ketolide antibiotic, or to any of the excipients.

Special warnings and precautions for use
Hypersensitivity
Rare cases of angioedema, serious allergic reactions including anaphylactic reactions and exceptionally rarely fatal outcomes have been reported with azithromycin. Some of these reactions with azithromycin have been recurrent in nature, requiring observation and treatment for a sufficient period of time. Symptoms may include facial edema, throat edema, chest tightness, dyspnea, urticaria and pruritus. Patients should be advised to discontinue use and seek medical attention immediately if they experience symptoms and required a longer period of observation and treatment.

Cardiac conduction
Prolongation of the QT interval has been reported with the use of macrolide antibiotics. This should be considered when azithromycin is prescribed to patients with known risk factors, including concomitant use with azithromycin.

Liver dysfunction
Cases of hepatic dysfunction, liver failure, and hepatitis have precipitated by co-administration of azithromycin and other medicinal products have been reported. Patients should be advised to discontinue use and seek medical attention immediately if they experience symptoms of liver dysfunction.

Diarrhoea
Mild diarrhoea is frequent, however severe diarrhoea requires discontinuation of azithromycin.

Cross-resistance
Azithromycin may induce resistance to macrolide antibiotics where their high dose concentrations may be used.

Clostridium difficile associated diarrhoea
Cases of *Clostridium difficile* associated diarrhoea (CDAD) have been reported with use of nearly all antibacterial agents, including azithromycin, and may range in severity from mild diarrhoea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhoea following the use of any antibacterial agent. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of such agents.

Renal impairment
In patients with severe renal impairment (GFR <10 mL/min), a 50% increase in systemic exposure (AUC) was observed. Dose adjustment is recommended in these patients.

Cardiovascular events
In a large, randomized, double-blind trial of azithromycin and CT interval, a trend in the risk of developing cardiovascular events (cardiovascular death, resuscitated cardiac arrest, and torsades de pointes) was observed in patients receiving azithromycin 500 mg daily for 5 days. Although the difference was not statistically significant, there was an increase in risk for cardiovascular events in patients receiving azithromycin. Patients should be advised to discontinue azithromycin and seek medical attention if they experience symptoms of cardiovascular events.

Myasthenia gravis exacerbation
Exacerbation of myasthenia gravis symptoms and new onset of myasthenia syndrome have been reported in patients receiving azithromycin therapy since market introduction.

Pyloric stenosis
Pyloric stenosis in neonates (prevention or treatment of *Mycobacterium avium* complex in children).

Lactose content
Zenith contains lactose, and effectiveness of tricyclines use of azithromycin in patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Neurological and psychiatric disorders
Azithromycin should be used with caution in patients with neurological and psychiatric disorders.

Interaction with other medicinal products and other forms of interaction
Antacids
Oral administration of azithromycin immediately after of simultaneous administration of antacid containing aluminium and magnesium hydroxides did not affect bioavailability as measured by serum concentrations. The medicinal products should not be taken concomitantly and significant changes in the GIT motility.

Cimetidine
Administration of cimetidine 2 hours before azithromycin did not affect azithromycin pharmacokinetic parameters.

Digoxin
Administration of azithromycin 0.5 g daily and azithromycin 250 mg daily did not alter the pharmacokinetic parameters of digoxin in healthy volunteers as determined by digoxin concentrations and renal clearance.

Ergot derivatives
Although pharmacokinetic interaction study in healthy volunteers, no significant effect was observed on the plasma levels of azithromycin or its tissue metabolite in patients receiving concomitant ergot derivatives.

Fluconazole
Co-administration of azithromycin 1200 mg and a single dose of 800 mg fluconazole had no significant effect on the pharmacokinetics of azithromycin. The overall exposure (AUC) and peak serum concentration (Cmax) of azithromycin were unaffected by fluconazole administration, and the observed changes in azithromycin half-life and oral clearance were not clinically significant.

Hexamethyldisilazane
Co-administration of azithromycin 1200 mg and efavirenz 400 mg daily for 7 days did not have significant effect on the pharmacokinetics of azithromycin.

Indinavir
Co-administration of azithromycin 1200 mg and indinavir 800 mg every 8 hours for 5 days did not significantly affect the pharmacokinetics of azithromycin.

Nelfinavir
Co-administration of azithromycin 1200 mg and nelfinavir at steady state 750 mg three times daily for 7 days resulted in changes in azithromycin concentrations that are not clinically significant. No dose adjustment is required.

Rifabutin
Co-administration of azithromycin and rifabutin did not affect the serum concentrations of either medicinal product.

Rifampicin
Co-administration of rifampicin to patients receiving concomitant treatment of azithromycin and rifampicin resulted in decreased serum concentrations of azithromycin. However, the clinical significance of this relationship is unknown as azithromycin has not been established clinical trials.

Sildenafil
Co-administration of azithromycin 500 mg daily for 3 days and sildenafil 100 mg did not affect the Cmax and AUC of sildenafil or its major circulating metabolite.

Zidovudine
Co-administration of azithromycin 1 g and zidovudine 600 mg did not significantly affect the pharmacokinetic parameters of zidovudine or its glucuronide metabolite. However, there have been case reports where the possibility of such an interaction between azithromycin and zidovudine could not be excluded.

Cetirizine
Co-administration of a clinically significant pharmacokinetic interaction when azithromycin was given with cetirizine.

Didanosine
Co-administration of azithromycin at concentrations of 500 mg for 3 days did not result in significant changes in didanosine plasma concentrations in comparison to placebo.

Fosamprenavir/ritonavir
Co-administration of azithromycin 1200 mg/day and fosamprenavir 700 mg/ritonavir 100 mg twice daily for 7 days did not significantly affect the pharmacokinetic parameters of azithromycin. However, azithromycin concentrations were similar to those seen in other studies.

Fluconazole/sulfamethoxazole/trimethoprim
Co-administration of azithromycin 1200 mg/day and fluconazole 800 mg/day for 7 days or with sulfamethoxazole/trimethoprim 800 mg/160 mg for 7 days did not significantly affect the pharmacokinetic parameters of azithromycin. However, azithromycin concentrations were similar to those seen in other studies.

Carbamazepine
In vitro studies, adequate data from the use of azithromycin in pregnant women. An epidemiological study suggested an increased risk of congenital malformations, but subsequent data have not confirmed this finding. Animal studies have shown reproductive toxicity. Zenith should not be used during pregnancy unless the benefit to the mother clearly outweighs any potential risk to the foetus. Azithromycin should only be used during pregnancy if the benefit outweighs the risk.

Lactation
Azithromycin is excreted in human milk. Caution should be exercised when Zenith is administered to a nursing woman. A decision should be made whether to discontinue nursing or to discontinue the medicinal product, taking into account the benefit of breastfeeding for the child and the benefit of therapy for the woman.

Fertility
Studies conducted in rat showed no adverse effects on fertility.

Undesirable effects
Adverse reactions possibly or probably related to Mycobacterium avium complex treatment in clinical trials or reported since marketing. Adverse reactions are ranked under headings of decreasing frequency: very common, common, uncommon, rare, very rare, not known, including isolated reports. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Gastrointestinal disorders
Very common: Diarrhoea, nausea, vomiting, abdominal pain, nausea, flatulence, abdominal discomfort, dyspepsia.
Uncommon: Vomiting, impaired, flatulence.

Hepatobiliary disorders
Uncommon: Hepatic function abnormal.

Skin and subcutaneous tissue disorders
Uncommon: Rash, pruritus.
Rare: Photosensitivity reaction, pruritus, photosensitivity reaction.

Musculoskeletal and connective tissue disorders
Uncommon: Joint pain.

Central and peripheral nervous system disorders
Uncommon: Dizziness, headache.
Rare: Asthenia, malaise.

Overdose
Oral administration of azithromycin to experimental animals does was similar to those seen at therapeutic doses. Animals received doses of azithromycin 200 mg/kg showed reversible loss of hearing, severe nausea, vomiting and diarrhoea.

Overdose in humans, including those children, has been associated with gastrointestinal symptoms. Treatment should be symptomatic and supportive as indicated.

PHARMACOLOGICAL PROPERTIES
Pharmacodynamic properties
Pharmacotherapeutic group: Macrolides for systemic use, macrolides, Azithromycin and ATC code: J01FA10.

Mechanism of action
Azithromycin is a macrolide of the macrolide subclass. By binding to the 50S ribosomal subunit, azithromycin arrests the translocation of peptidyl-tRNA from one site to the next site on the ribosome, thereby inhibiting protein synthesis.

PHARMACOKINETICS
The efficacy of azithromycin is best described by the relationship AUC/MIC, where AUC = 24-hour area under the concentration-time curve and MIC = minimum inhibitory concentration.

Absorption of azithromycin
Following oral administration, azithromycin is rapidly absorbed and is more acid stable. Peak serum concentrations are usually achieved within 2 to 3 hours. Absolute bioavailability is approximately 37%. Food does not affect the bioavailability of azithromycin. Maximum serum concentrations were observed 2 to 4 hours after oral administration.

Distribution of azithromycin
Azithromycin can be used to determine susceptibility to azithromycin, clarithromycin and erythromycin.

Azithromycin is widely distributed throughout the body following oral dosing. Measurement of tissue to plasma concentration ratios following multiple dosing of 500 mg daily for 3 days to healthy volunteers showed that the ratio of azithromycin in lung tissue to plasma was 25 to 1. Tissue concentrations of azithromycin were substantially higher than corresponding plasma concentrations, suggesting that the drug localizes in tissues. Plasma protein binding of azithromycin is variable, depending on plasma concentration and the type of tissue involved, and decreases as concentrations increase to 51% at a concentration of 0.02 mcg/mL.

Metabolism of azithromycin
Azithromycin undergoes N-demethylation in the liver and is chemically similar to erythromycin. Azithromycin is also metabolized by hydroxylation of the desosamine and aglycone rings. Azithromycin is not a potent inhibitor of the cytochrome P450 system in human liver microsomes. Azithromycin does not induce hepatic microsomal enzymes.

Elimination of azithromycin
Azithromycin is eliminated primarily in unchanged form in bile and is excreted in high concentrations as active drug in the urine over the 3 days following administration. Total clearance of azithromycin is high, with hepatic clearance approximating hepatic blood flow. Azithromycin undergoes enterohepatic recirculation. Terminal elimination half-life of azithromycin is 68 hours, due to extensive uptake into tissues and subsequent release from tissues. Azithromycin is excreted in the bile as unchanged drug and in the urine as unchanged drug and as metabolites.

Pharmacokinetics in special populations
Renal impairment
After oral administration, pharmacokinetic parameters in adult subjects (18 to 3 years) with normal renal function (creatinine clearance ≥80 mL/min) and in subjects with mild (creatinine clearance 40 to 80 mL/min) or moderate (creatinine clearance 10 to 40 mL/min) renal impairment who received a single 1 g dose of azithromycin. Peak serum concentrations (Cmax) and total exposure (AUC) were not significantly affected by renal impairment.

In a similar study, high concentrations of azithromycin were found in the kidneys of animals given azithromycin. Azithromycin is primarily excreted in bile in unchanged form. Renal clearance accounts for approximately 6% of the total body clearance of azithromycin. Dose adjustment in patients with renal impairment is not necessary.

Hepatic impairment
Azithromycin is primarily excreted unchanged in the liver. Patients with cirrhosis showed no increase in half-life of azithromycin.

Elderly
Pharmacokinetic parameters were not significantly affected in the therapeutic range.

Paediatrics
Pharmacokinetic studies of the concentrations of azithromycin measured in tissues are supported by microbiological data. Following oral administration of 10 mg/kg to children (aged 6 months to 12 years), the majority is excreted in the first 24 hours. The terminal half-life during this period is 68 hours. Clearance is approximately 80% of that in adults, resulting in similar exposure in children receiving 10 mg/kg and adults receiving 500 mg.

Pharmacodynamics in animal models of infection
Time-kill studies

Time-kill studies have demonstrated that azithromycin is concentration and time dependent in its killing of susceptible bacteria. The minimum bactericidal concentration (MBC) is usually equal to the minimum inhibitory concentration (MIC). Azithromycin is rapidly bactericidal against *Haemophilus influenzae* at concentrations equal to the MIC.

Post-antibiotic effect

Azithromycin exhibits a significant post-antibiotic effect against *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Streptococcus pyogenes* and *Legionella pneumophila*. The post-antibiotic effect persists for 1.2 hours after exposure to azithromycin at concentrations equal to the MIC.

Efficacy in animal models of infection

Azithromycin has been active in animal models of infection with *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Streptococcus pyogenes* and *Legionella pneumophila*.

Clinical efficacy

Azithromycin has been active in clinical infections caused by *Haemophilus influenzae*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, *Streptococcus pyogenes* and *Legionella pneumophila*.

Pharmaceutical particulars
Incompatibilities
None known.

STORAGE
Store at temperature not exceeding 30°C.

CAUTION
Keep out of reach of children. Dangerous if swallowed. Avoid prohibited dispensing without prescription.

AVAILABILITY
Blister packs of 3 tablets (10's).

ADVERSE DRUG REACTION REPORTING STATEMENT
Adverse drug reactions should be reported to the Bureau of Food and Drugs, Department of Health, Sta. Escolastica Compound, San Lazaro Hospital, Quezon City, Metro Manila, Philippines. Tel. No. 7-1-65-700 to 09. Fax No. 711-9412. E-mail: postmaster@fda.doh.gov.ph.

DATE OF FIRST AUTHORIZATION / REGISTRATION NUMBER
DOH-FDR-0000-14

DATE OF REVISION
January 27, 2016

RETAIL PRICE NOT TO EXCEED Php. 131.43
WHOSE DRUG PRICE REGULATION

UNILAB Consumer Care
8-UNILAB-1
8-8645-2251
Philippines: 1-800-1-8-UNILAB-1
International: +63-2-8-8645-2251
Website: www.unilab.com.ph

Manufactured by: Mercantile Bangladesh Limited
Tongi Plant, Gazaria, Bl. Mymen, Tongi, 1711 Gazipur, Bangladesh
Distributed by: United Laboratories, Inc. (Unilab)
Mandaluyong City, Philippines

13-10-2276/R03
13-10-2276/R03