UNILAB

Acetylcysteine

Exflem

200 mg Powder for Oral Solution

MUCOLYTIC

℞

FORMULATION

Each sachet contains:

Acetylcysteine ..... 200 mg

PHARMACEUTICAL FORM

Powder for oral solution

CLINICAL PARTICULARS

Therapeutic Indications

Secretolytic therapy in acute and chronic bronchopulmonary diseases accompanied by impaired formation and transport of mucus.

DOSAGE AND METHOD OF ADMINISTRATION

If not otherwise prescribed, the following dosage is recommended for acetylcysteine:

Adults and adolescents from 14 years of age

1 powder sachet 2-3 times daily (equivalent to 400 - 600 mg acetylcysteine/day)

Children and adolescents 6-14 years of age

1 powder sachet twice daily (equivalent to 400 mg acetylcysteine/day)

Method of administration

Acetylcysteine is to be taken dissolved in a glass of water after meals.

Duration of use

Acetylcysteine should not be taken for more than 4-5 days without medical advice.

In case of chronic bronchitis and mucociliaric insufficiency, long-term treatment should be performed in order to achieve effective prophylaxis.

CONTRAINDICATIONS

- Hypersensitivity to acetylcysteine or to any of the excipients
- Active peptic ulceration
- Children below 2 years of age

SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Patients with bronchial asthma must be closely monitored during therapy. If bronchospasm occurs, the use of acetylcysteine must be discontinued immediately and appropriate treatment initiated.

Caution is advised when using this product in patients with a history of peptic ulcer, especially if additional drugs are being taken that are known to irritate the mucous membranes of the gastrointestinal tract.

The use of acetylcysteine, especially in early treatment can lead to liquefaction and thus to an increase in volume of bronchial secretions. If the patient is unable to cough (sufficiently expectorate), appropriate measures (such as drainage and aspiration) should be taken.

The occurrence of severe skin reactions such as Stevens-Johnson syndrome and Lyell's syndrome has very rarely been reported in temporal connection with the use of acetylcysteine. If symptoms or signs of severe skin lesions, newly occur, medical advice should be sought without delay and use of acetylcysteine be terminated (see also Undesirable effects).

Caution is advised in patients with histamine intolerance. Treatment with acetylcysteine for longer periods should be avoided in such patients, as acetylcysteine affects histamine metabolism and may lead to symptoms of intolerance (e.g. headache, runny nose, itching).

Children and adolescents

Mucolytics can result in blockage of the respiratory tract in children under 2 years of age. They are not able to expectorate their respiratory tract and their limited ability to cough up mucus. Therefore, mucolytics must not be used in children under 2 years of age (see Contraindications).

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

Combined administration of acetylcysteine with antitussives may cause a dangerous secretory congestion due to the elevated cough reflex, so that an especially careful diagnosis is required for this combined treatment.

Reports to date on an inactivation of antibiotics due to acetylcysteine exclusively refer to in vitro experiments in which the relevant substances were mixed in one vessel. Nevertheless for safety reasons, oral antibiotics should be administered separately and at an interval of at least 2 hours. This does not apply to cefixime and broad-spectrum penicillins.

Acetylcysteine/glyceryl trinitrate

The concomitant administration of acetylcysteine can potentially result in an intensification of the vasodilatory and inhibition of platelet aggregation effects of glyceryl trinitrate (nitroglycerine).

Concomitant treatment with glyceryl trinitrate and acetylcysteine is considered necessary, patients should be monitored for a possible development of increased headache, which can be serious, and advised of the possibility of headaches.

Activated carbon in high doses (as an antidote) can reduce the effectiveness of acetylcysteine.

Changes in the determination of laboratory parameters

Acetylcysteine can affect the colorimetric determination of salicylates.

In urine tests, acetylcysteine can affect the results of determinations of ketone bodies.

The dissolution of acetylcysteine formulations together with other medicinal products is not recommended.

Pregnancy and lactation

Pregnancy

No sufficient data on exposed pregnant women are available for acetylcysteine. Experimental animal studies do not suggest direct or indirect harmful effects on pregnancy, embryonal/fetal development, parturition or postnatal development (see also Preclinical safety data). Acetylcysteine should be used during pregnancy after strict assessment of the benefit/risk ratio.

Lactation

No information is available regarding excretion into breast milk. Acetylcysteine should be used during lactation only after strict assessment of the benefit/risk ratio.

Fertility

No data are available on the effect of acetylcysteine on human fertility. In animal studies, no adverse effects on fertility were observed at therapeutic doses of acetylcysteine (see Preclinical safety data).

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Acetylcysteine has no known effect on the ability to drive and use machines.

UNDESIRABLE EFFECTS

The evaluation of adverse reactions is based on the following information on frequencies:

Very common: (≥ 1/10)

Common: (≥ 1/100 up to < 1/10)

Uncommon: (≥ 1/1,000 up to < 1/100)

Rare: (≥ 1/10,000 up to < 1/1,000)

Very rare: (< 1/10,000)

Not known: (cannot be estimated from the available data)

Immune system disorders

Uncommon: Hypersensitivity reactions

Very rare: Anaphylactic shock, anaphylactic/ anaphylactoid reactions

Nervous system disorders

Uncommon: Headache

Ear and labyrinth disorders

Uncommon: Tinnitus

Cardiac disorders

Uncommon: Tachycardia

Vascular disorders

Very rare: Hemorrhage

Respiratory, thoracic and mediastinal disorders

Rare: Dyspnoea, bronchospasm

Gastrointestinal disorders

Uncommon: Nausea, vomiting, diarrhoea, abdominal pain, stomatitis

Rare: Dyspepsia

Skin and subcutaneous tissue disorders

Uncommon: Urticaria, rash, angioedema, itching, exanthema

General disorders and administration site conditions

Uncommon: Fever

Not known: Facial edema

Investigations

Uncommon: Hypotension

In very rare cases, severe skin reactions such as Stevens-Johnson syndrome and toxic epidermal necrolysis have been reported in temporal connection with the use of acetylcysteine. As most of these reports contain other, at least one additional drug that could potentially have intensified the described mucocutaneous effects was being taken at the same time.

If skin or mucous membrane abnormalities develop, medical advice should therefore immediately be sought and the use of acetylcysteine discontinued.

A decreased blood platelet aggregation in the presence of acetylcysteine has been confirmed by different studies. The clinical relevance has not yet been clarified to date.

Overdose

Symptoms of toxic overdose has been observed to date in association with cell phenomena and forms of acetylcysteine. Volunteers were treated with a dose of 11.8 g acetylcysteine/kg body weight 3 months without observing any severe side effects. Oral doses up to 500 mg acetylcysteine/kg BW were tolerated without any symptoms of intoxication.

Symptoms of intoxication

Overdoses may lead to gastrointestinal symptoms, such as nausea, vomiting and diarrhoea. Infants are at risk of hypersecretion.

Therapy of intoxication

If necessary, according to the symptoms.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Pharmacotherapeutic group: mucolytics

ATC Code: R05CB01

Acetylcysteine is a derivative of the amino acid cysteine. The efficacy of acetylcysteine is secretolytic and secretomotoric in the area of the respiratory tract. It is discussed that it splits off the interconnecting disulphide bonds between the mucin molecules and thus that it also has a depolymerising effect on DNA-chains (in purulent mucus). Due to these mechanisms, the mucus viscosity should be reduced.

An alternative explanation of acetylcysteine is meant to be based on the capacity of its reactive SH group to bind chemical radicals and to detoxify them in this way.

Furthermore, acetylcysteine contributes to an increase in glutathione synthesis, which is decreased in the case of detoxification of noxae. This provides the explanation for its antidotal effect in paracetamol intoxication.

Pharmacokinetic properties

Absorption

Following oral administration, acetylcysteine is rapidly and almost completely absorbed and metabolised in the liver (first-pass effect). Peak plasma levels are reached, as well as to deacetylcysteine, cystine and further mixed disulfides.

Distribution

Due to the high first-pass effect, the bioavailability of orally administered acetylcysteine is very low (approx. 10%). The plasma protein binding of acetylcysteine is 50% and after 1.5 hours with the maximum plasma concentration of the metabolite cysteine in the range of approx. 2 mmol/L. The protein binding of acetylcysteine was determined to be about 50%.

Biotransformation

The acetylcysteine and its metabolites occur in three different forms in the organism: partially in free form, as a disulfide bound to proteins via labile disulphide bonds and partially as incorporated amino acid. Acetylcysteine is excreted almost exclusively in the form of inactive metabolites, inorganic sulphates, deacetylcysteine and neutral sulphur.

Elimination

Pharmacokinetic studies with intravenous administration of acetylcysteine revealed a distribution volume of 0.47 L/kg and a total clearance of 0.11 L/h/kg (in total) and 0.34 L/h/kg (renal), respectively.

No knowledge is available concerning the behaviour of acetylcysteine at the blood-brain barrier in humans.

Preclinical safety data

Acute toxicity

There is low toxicity in animal experiments is low. For the treatment of overdoses, see Overdose.

Chronic toxicity

Studies in various animal species (rat, dog) with a duration of up to one year did not show any pathological alterations.

Teratogenic and mutagenic potential

No mutagenic effects of acetylcysteine are to be expected. An in vivo test was negative.

No studies of a teratogenic potential of acetylcysteine have been carried out.

Reproductive toxicology

No malformations were detected in embryotoxicity studies in rabbits and rats. Studies of fertility and perinatal or postnatal toxicity were negative.

Acetylcysteine passes the placenta in rats and was detected in the foetal tissue. Administration of the metabolite L-cysteine to rats led to a maximum plasma concentration in placenta and foetus for up to 8 hours after oral administration.

PHARMACEUTICAL PARTICULARS

Incompatibilities

Not applicable

ADVERSE DRUG REACTION REPORTING STATEMENT

For suspected adverse drug reaction, seek medical attention immediately and report to the FDA at www.fda.gov.ph/adr/ and/or +632-8-6944888 / +632-8-8443522 / Dr. Mario Mondia Hotline +1-800-10UNILAB / for provinces, e-mail pvg@unilab.com.ph.

By reporting undesirable effects, you can help provide more information on the safety of this medicine.

CAUTION

Foods, Drugs, Devices, and Cosmetics Act prohibits dispensing without a prescription.

STORAGE CONDITION

Store at temperatures not exceeding 30°C.

AVAILABILITY

Thermosealed Triplefoyer Polyethylene Aluminum paper sachet x 2's Box of 20's

Date of First Authorization: May 2018

Date of Revision: May 2020; COS-05

DRT-7&31

The elimination halflife after intravenous administration is 30-40 minutes while excretion follows threephase kinetics (alpha, beta, and terminal gamma phase).

Acetylcysteine crosses the placenta and is detected in cord blood. No information is available regarding excretion into breast milk.

The elimination halflife after intravenous administration is 30-40 minutes while excretion follows threephase kinetics (alpha, beta, and terminal gamma phase).

Acetylcysteine crosses the placenta and is detected in fetal organs. No information is available regarding excretion into breast milk.

Manufactured by: Laboratorios Grin®
Noucella #2, 31690 VT Villava, Germany
Under authority by: Unilab Inc.
Bridgeway Corporate Center, Alabang, Muntinlupa
Laboratorios Grin Philippine Operations Office
Unit 1, 2nd Floor, Building 1, Cyberpod Centris, Quezon City
Apollo Asia Corp. - Edificio and Amorsolo Sts., Brgy. San Lorenzo, Makati City
Tel. No. 8-8443522 / 8-8443523
Via: 6th United Street, Mandaluyong City
Metro Manila, Philippines

UNILAB Customer Care
0-UNILAB-1
0-864-522-1
Mon-Fri 8:00AM - 5:00PM
Email: consumer@unilab.com.ph
www.unilab.com.ph

Reg. #PCPHL
50086706
PED15715ONN01
40131111